

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Docket No.: PF-0379-1 DIV

Response Under 37 C.F.R. 1.116 - Expedited Procedure
Examining Group 1652

Certificate of Mailing

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to:
Mail Stop Appeal Brief-Patents, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450 on _____.

By: _____

Printed: _____

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
BEFORE THE BOARD OF PATENT APPEALS AND INTERFERENCES**

In re Application of: Bandman et al.

Title: HUMAN MITOCHONDRIAL MALATE DEHYDROGENASE

Serial No.: 09/915,694

Filing Date:

July 25, 2001

Examiner: Fronda, C.

Group Art Unit:

1652

Mail Stop Appeal Brief-Patents
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

BRIEF ON APPEAL

Sir:

Further to the Notice of Appeal filed on September 30, 2003, and received by the USPTO on October 6, 2003, herewith are three copies of Appellants' Brief on Appeal. Appellants hereby request a one month extension of time in order to file this Brief. Authorized fees include the statutory fee of \$110 for a one-month extension of time, as well as the \$ 330.00 fee for the filing of this Brief.

This is an appeal from the decision of the Examiner finally rejecting claims 3, 6, 7, 9 and 12 of the above-identified application.

01/13/2004 MGBREM1 00000034 090108 09915694

01 FC:1402 330.00 DA

(1) REAL PARTY IN INTEREST

The above-identified application is assigned of record to Incyte Pharmaceuticals, Inc. (now Incyte Corporation, formerly known as Incyte Genomics, Inc.) (Reel 9774, Frame 0975), which is the real party in interest herein.

(2) RELATED APPEALS AND INTERFERENCES

Appellants, their legal representative and the assignee are not aware of any related appeals or interferences which will directly affect or be directly affected by or have a bearing on the Board's decision in the instant appeal.

(3) STATUS OF THE CLAIMS

Claims rejected: Claims 3, 6, 7, 9 and 12
Claims allowed: (none)
Claims objected to: Claims 4, 5 and 10.
Claims canceled: Claims 1, 2, 8, 11, 13, 17-27, 30-45.
Claims withdrawn: Claims 14-16, 28 and 29
Claims on Appeal: Claims 3, 6, 7, 9 and 12 (A copy of the claims on appeal, as amended, can be found in the attached Appendix).

(4) STATUS OF AMENDMENTS AFTER FINAL

There were no amendments made after final.

(5) SUMMARY OF THE INVENTION

Embodiments of the present invention are directed, *inter alia*, to polynucleotides encoding mitochondrial malate dehydrogenase (MT-MDH). In particular embodiments, the mitochondrial malate dehydrogenases are selected from among amino acid sequences comprising SEQ ID NO:1. These polypeptides have strong chemical and structural homology with known mitochondrial malate dehydrogenases. For example:

MT-MDH is 338 amino acids in length and has two potential N-glycosylation sites at residues N-117 and N-145, seven potential casein kinase II phosphorylation sites at T-54, S-69, T-109, T-170, S-261, S-309, and S-310, four potential protein kinase C phosphorylation sites at residues T-213, T-227, S-326, and T-336, a mitochondrial malate dehydrogenase active site signature between residues V-169 and V-181, and a transit peptide sequence from residues M-1 to N-24. As shown in Figures 2A and 2B, MT-MDH has chemical and structural homology with murine mitochondrial mitochondrial malate dehydrogenase (GI 56643; SEQ ID NO:3) and porcine mitochondrial mitochondrial malate dehydrogenase (GI 164541; SEQ ID NO:4). In particular, MT-MDH and murine

mitochondrial mitochondrial malate dehydrogenase share 94% identity, share both potential N-glycosylation sites, six potential casein kinase II sites, three potential protein kinase C sites, the mitochondrial malate dehydrogenase active site signature, and the transit peptide sequence. As illustrated by Figures 3A and 3B, respectively, MT-MDH and murine mitochondrial mitochondrial malate dehydrogenase (SEQ ID NO:3) have similar isoelectric points (pI = 8.8). As illustrated by Figures 4A and 4B, MT-MDH contains potential NAD(H) and NADP(H) binding site motifs. Northern analysis shows the expression of this sequence in various libraries, at least 49% of which are immortalized or cancerous and at least 24% of which involve immune response. Of particular note is the expression of MT-MDH in fetal tissues; in cardiovascular, gut, nervous, and reproductive tissues; and in secretory and hematopoietic tissues. (Specification at page 14, line 29 to page 15, line 17).

The polynucleotides of the present invention have a variety of utilities. For example, they can be used for the diagnosis, prevention, or treatment of vesicle trafficking, immunological and neoplastic disorders. (See the Specification e.g., at page 14, lines 18-21)

(6) ISSUES

1. Whether claims 3, 6, 7, 9 and 12 meet the written description requirement of 35 U.S.C. § 112, first paragraph.
2. Whether claims 3, 6, 7, 9 and 12 meet the enablement requirement of 35 U.S.C. § 112, first paragraph.

(7) GROUPING OF THE CLAIMS

As to Issue 1

All of the claims on appeal are grouped together.

As to Issue 2

All of the claims on appeal are grouped together.

(8) APPELLANTS' ARGUMENTS

Issue 1- Written description rejection under 35 U.S.C. § 112, first paragraph

Claims 3, 6, 7, 9 and 12 were rejected under 35 U.S.C. §112, first paragraph, as allegedly

“containing subject matter which was not described in the specification in such a way as to reasonably convey to one skilled in the relevant art the inventor(s) at the time the application was filed, had possession of the claimed invention.” (06/20/03 Office Action, at page 4) In making the rejection, the Examiner asserts that:

There is no disclosure of any particular structure to function/activity relationship in the single disclosed species. The specification also fails to described additional representative species of these polynucleotides by any identifying structural characteristics or properties for which no predictability of structure is apparent. Given this lack of additional representative species as encompassed by the claims, Applicants have failed to sufficiently disclose the invention in such full, clear, concise and exact terms that a skilled artisan would recognize Applicants were in possession of the claimed invention. (12/17/02 Office Action, at pages 4-5).

This rejection is improper, as the claims define subject matter which is described in the Specification in such a way as to reasonably convey to one skilled in the art that the inventors had possession of the claimed subject matter at the time the application was filed. The requirements necessary to fulfill the written description requirement of 35 U.S.C. § 112, first paragraph, are well established by case law:

... the applicant must also convey with reasonable clarity to those skilled in the art that, as of the filing date sought, he or she was in possession of *the invention*. The invention is, for purposes of the “written description” inquiry, *whatever is now claimed*. *Vas-Cath, Inc. v. Mahurkar*, 19 U.S.P.Q.2d 1111, 1117 (Fed. Cir. 1991).

The Board’s attention is also drawn to the Patent and Trademark Office’s own Examination of Patent Applications Under the 35 U.S.C. § 112, para. 1”, published January 5, 2001, which provide that:

An applicant may also show that an invention is complete by disclosure of sufficiently detailed, relevant identifying characteristics⁴² which provide evidence that applicant was in possession of the claimed invention,⁴³ i.e., complete or partial structure, other physical and/or chemical properties, functional characteristics when coupled with a known or disclosed correlation between function and structure, or some combination of such characteristics.⁴⁴ What is conventional or well known to one of ordinary skill in the art need not be disclosed in detail.⁴⁵ If a skilled artisan would have understood the inventor to be in possession of the claimed invention at the time of filing, even if every nuance of

the claims is not explicitly described in the specification, then the adequate description requirement is met.⁴⁶

Thus, the written description standard is fulfilled by both what is specifically disclosed and what is conventional or well known to one skilled in the art.

A. The specification provides an adequate written description of the claimed “variants” of SEQ ID NO:1 and SEQ ID NO:2.

The subject matter encompassed by claims 3, 6, 7, 9 and 12 is either disclosed by the specification or is conventional or well known to one skilled in the art.

First note that the "variant" language of independent claim 3 recites a polynucleotide encoding "a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1" and the "variant" language of independent claim 12 recites "a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2."

The amino acid sequence of SEQ ID NO:1 and the polynucleotide sequence of SEQ ID NO:2 are explicitly disclosed in the specification. See, for example, the Sequence Listing. Variants of SEQ ID NO:1 and SEQ ID NO:2 are described in the Specification at, for example, page 4, lines 13-14; page 7, lines 4-7 and 12-18; and page 15, lines 18-27.

One of ordinary skill in the art would recognize polynucleotide sequences which are variants having a polynucleotide sequence at least 95% to SEQ ID NO:2, or which encode polypeptide variants having an amino acid sequence at least 95% identical to SEQ ID NO:1. Given any naturally occurring polynucleotide sequence, it would be routine for one of skill in the art to recognize whether it was a variant of SEQ ID NO:2, or whether it encoded a variant of SEQ ID NO:1. Accordingly, the specification provides an adequate written description of the recited polynucleotide variants of SEQ ID NO:2 and polynucleotides encoding polypeptide variants of SEQ ID NO:1.

1. The present claims specifically define the claimed genus through the recitation of chemical structure

Court cases in which “DNA claims” have been at issue commonly emphasize that the

recitation of structural features or chemical or physical properties are important factors to consider in a written description analysis of such claims. For example, in *Fiers v. Revel*, 25 U.S.P.Q.2d 1601, 1606 (Fed. Cir. 1993), the court stated that:

If a conception of a DNA requires a precise definition, such as by structure, formula, chemical name or physical properties, as we have held, then a description also requires that degree of specificity.

In a number of instances in which claims to DNA have been found invalid, the courts have noted that the claims attempted to define the claimed DNA in terms of functional characteristics without any reference to structural features. As set forth by the court in *University of California v. Eli Lilly and Co.*, 43 U.S.P.Q.2d 1398, 1406 (Fed. Cir. 1997):

In claims to genetic material, however, a generic statement such as “vertebrate insulin cDNA” or “mammalian insulin cDNA,” without more, is not an adequate written description of the genus because it does not distinguish the claimed genus from others, except by function.

Thus, the mere recitation of functional characteristics of a DNA, without the definition of structural features, has been a common basis by which courts have found invalid claims to DNA. For example, in *Lilly*, 43 U.S.P.Q.2d at 1407, the court found invalid for violation of the written description requirement the following claim of U.S. Patent No. 4,652,525:

1. A recombinant plasmid replicable in procaryotic host containing within its nucleotide sequence a subsequence having the structure of the reverse transcript of an mRNA of a vertebrate, which mRNA encodes insulin.

In *Fiers*, 25 U.S.P.Q.2d at 1603, the parties were in an interference involving the following count:

A DNA which consists essentially of a DNA which codes for a human fibroblast interferon-beta polypeptide.

Party Revel in the *Fiers* case argued that its foreign priority application contained an adequate written description of the DNA of the count because that application mentioned a potential method for isolating the DNA. The Revel priority application, however, did not have a description of any particular DNA structure corresponding to the DNA of the count. The court therefore found that the Revel priority application lacked an adequate written description of the

subject matter of the count.

Thus, in *Lilly* and *Fiers*, nucleic acids were defined on the basis of functional characteristics and were found not to comply with the written description requirement of 35 U.S.C. §112; *i.e.*, "an mRNA of a vertebrate, which mRNA encodes insulin" in *Lilly*, and "DNA which codes for a human fibroblast interferon-beta polypeptide" in *Fiers*. In contrast to the situation in *Lilly* and *Fiers*, the claims at issue in the present application define polynucleotides and polypeptides in terms of chemical structure, rather than on functional characteristics. For example, the "variant language" of independent claims 3 and 12 recite chemical structure to define the claimed genus:

3. An isolated polynucleotide encoding a polypeptide selected from the group consisting of:.....

b) a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.

12. An isolated polynucleotide selected from the group consisting of:

b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2,....

From the above it should be apparent that the claims of the subject application are fundamentally different from those found invalid in *Lilly* and *Fiers*. The subject matter of the present claims is defined in terms of the chemical structure of SEQ ID NO:1 and SEQ ID NO:2. In the present case, there is no reliance merely on a description of functional characteristics of the polynucleotides and polypeptides recited by the claims. In fact, there is no recitation of functional characteristics. Moreover, if such functional recitations were included, it would add to the structural characterization of the recited polynucleotides. The polynucleotides defined in the claims of the present application recite structural features, and cases such as *Lilly* and *Fiers* stress that the recitation of structure is an important factor to consider in a written description analysis of claims of this type. By failing to base its written description inquiry "on whatever is now claimed," the Office Action fails to provide an appropriate analysis of the present claims and how they differ from those found not to satisfy the written description requirement in *Lilly* and *Fiers*.

2. The present claims do not define a genus which is "highly variant"

Furthermore, the claims at issue do not describe a genus which could be characterized as "highly variant." Available evidence illustrates that the claimed genus is of narrow scope.

In support of this assertion, the Board's attention is directed to the enclosed reference by Brenner et al. ("Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships," Proc. Natl. Acad. Sci. USA (1998) 95:6073-6078). Through exhaustive analysis of a data set of proteins with known structural and functional relationships and with <90% overall sequence identity, Brenner et al. have determined that 30% identity is a reliable threshold for establishing evolutionary homology between two sequences aligned over at least 150 residues. (Brenner et al., pages 6073 and 6076.) Furthermore, local identity is particularly important in this case for assessing the significance of the alignments, as Brenner et al. further report that $\geq 40\%$ identity over at least 70 residues is reliable in signifying homology between proteins. (Brenner et al., page 6076.)

The present application is directed, *inter alia*, to polynucleotides encoding mitochondrial malate dehydrogenases, including polynucleotides encoding mitochondrial malate dehydrogenases related to the amino acid sequence of SEQ ID NO:1. In accordance with Brenner et al, naturally occurring molecules may exist which could be characterized as mitochondrial malate dehydrogenases and which have as little as 30% identity over at least 150 residues to SEQ ID NO:1. The "variant language" of the present claims recites, for example, polynucleotides encoding a polypeptide comprising "a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1" (note that SEQ ID NO:1 has 338 amino acid residues). This variation is far less than that of polynucleotides encoding all potential mitochondrial malate dehydrogenases related to SEQ ID NO:1, i.e., those mitochondrial malate dehydrogenases having as little as 30% identity over at least 150 residues to SEQ ID NO:1.

3. The state of the art at the time of the present invention is further advanced than at the time of the *Lilly* and *Fiers* applications

In the *Lilly* case, claims of U.S. Patent No. 4,652,525 were found invalid for failing to comply with the written description requirement of 35 U.S.C. §112. The '525 patent claimed the

benefit of priority of two applications, Application Serial No. 801,343 filed May 27, 1977, and Application Serial No. 805,023 filed June 9, 1977. In the *Fiers* case, party Revel claimed the benefit of priority of an Israeli application filed on November 21, 1979. Thus, the written description inquiry in those case was based on the state of the art at essentially at the "dark ages" of recombinant DNA technology.

The present application has a priority date of September 3, 1997. Much has happened in the development of recombinant DNA technology in the 18 or more years from the time of filing of the applications involved in *Lilly* and *Fiers* and the present application. For example, the technique of polymerase chain reaction (PCR) was invented. Highly efficient cloning and DNA sequencing technology has been developed. Large databases of protein and nucleotide sequences have been compiled. Much of the raw material of the human and other genomes has been sequenced. With these remarkable advances one of skill in the art would recognize that, given the sequence information of SEQ ID NO:1 and SEQ ID NO:2, and the additional extensive detail provided by the subject application, the present inventors were in possession of the claimed polynucleotide variants at the time of filing of this application.

4. Summary

The 12/17/02 and 6/30/03 Office Actions fail to base the written description inquiry "on whatever is now claimed." Consequently, the Office Actions do not provide an appropriate analysis of the present claims and how they differ from those found not to satisfy the written description requirement in cases such as *Lilly* and *Fiers*. In particular, the claims of the subject application are fundamentally different from those found invalid in *Lilly* and *Fiers*. The subject matter of the present claims is defined in terms of the chemical structure of SEQ ID NO:1 and SEQ ID NO:2. The courts have stressed that structural features are important factors to consider in a written description analysis of claims to nucleic acids and proteins. In addition, the genus of polynucleotides defined by the present claims is adequately described, as evidenced by Brenner et al. Furthermore, there have been remarkable advances in the state of the art since the *Lilly* and *Fiers* cases, and these advances were given no consideration whatsoever in the position set forth by the Office Action.

For at least the reasons set forth above, the specification provides an adequate written description of the claimed subject matter, and this rejection should be reversed.

Issue 2 – Enablement rejection under 35 U.S.C. § 112, first paragraph

Claims 3, 6, 7, 9 and 12 stand rejected under 35 U.S.C. 112, first paragraph allegedly for lacking an enabling disclosure with respect to variants of SEQ ID NO:1. The Examiner has specifically stated that these claims contain “subject matter which was not described in the specification in such a way as to enable one skilled in the art to which it pertains, or with which it is most nearly connected, to make and/or use the invention.” (06/30/03 Office Action, at page 2-3)

In making the rejection, the Examiner asserts that:

“The specification does not provide guidance with respect to the specific structural/catalytic amino acids and the structural motifs essential for enzyme structure and activity function which cannot be altered. Thus searching for the specific nucleotides to change (deletion, insertion, substitution, or combinations thereof) in a polynucleotide is well outside the realm of routine experimentation and predictability in the art” (12/17/02 Office Action, at page 6).

The Examiner makes a similar assertion in the 06/30/03 Office Action that “[t]he specification does not teach the specific amino acids that can be altered and yet still retain enzyme activity.” (6/30/03 Office Action, at page 3).

The first paragraph of 35 U.S.C. §112 requires that the Specification describe how to make and use the claimed subject matter. That requirement has been met in the present application. In particular, the Specification describes how to make and use naturally-occurring polypeptide variants of SEQ ID NO:1 and polynucleotides encoding such variants.

Independent claim 3 recites not only that the “variant” polynucleotides encode polypeptides that are at least 95% identical to SEQ ID NO:1, but also have “*a naturally-occurring amino acid sequence.*” Through the process of natural selection, nature will have determined the appropriate amino acid sequences. Given the information provided by SEQ ID NO:1 (the amino acid sequence of MT-MDH) and SEQ ID NO:2 (the polynucleotide sequence encoding MT-MDH), one of skill in the art would be able to routinely obtain a polynucleotide encoding a polypeptide comprising “a naturally-occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.” Likewise for the “variant” polynucleotides defined by independent claim 12: “a polynucleotide comprising a naturally

occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2.” For example, the identification of relevant polynucleotides could be performed by hybridization and/or PCR techniques that were well-known to those skilled in the art at the time the subject application was filed and/or described throughout the Specification of the instant application. For example:

The terms “stringent conditions” or “stringency”, as used herein, refer to the conditions for hybridization as defined by the nucleic acid, salt, and temperature. These conditions are well known in the art and may be altered in order to identify or detect identical or related polynucleotide sequences. Numerous equivalent conditions comprising either low or high stringency depend on factors such as the length and nature of the sequence (DNA, RNA, base composition), nature of the target (DNA, RNA, base composition), milieu (in solution or immobilized on a solid substrate), concentration of salts and other components (e.g., formamide, dextran sulfate and/or polyethylene glycol), and temperature of the reactions (within a range from about 5°C below the melting temperature of the probe to about 20°C to 25°C below the melting temperature). One or more factors may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions. (Specification at page 13, lines 11-21)

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding MT-MDH or closely related molecules, may be used to identify nucleic acid sequences which encode MT-MDH. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding MT-MDH, alleles, or related sequences. (Specification at page 37, line 25 to page 38, line 3)

Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the MT-MDH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring MT-MDH. (Specification at page 38, lines 4-8)

See also Example VI, at page 50.

Thus, one skilled in the art need not make and test vast numbers of polypeptides that are based on the amino acid sequence of SEQ ID NO:1. Instead, one skilled in the art need only screen a cDNA library or use appropriate PCR conditions to identify relevant polynucleotides/polypeptides that already exist in nature. By adjusting the nature of the probe or nucleic acid (*i.e.*, non-conserved, conserved or highly conserved) and the conditions of hybridization (maximum, high, intermediate or low stringency), one can obtain variant polynucleotides of SEQ ID NO:2 which, in turn, will allow one to make the variant polypeptides of SEQ ID NO:1 recited by the present claims. Furthermore, the Specification sets forth an assay for measuring malate dehydrogenase activity (Example X at page 52, lines 11-18).

Accordingly, the document cited by the Examiner in the 12/17/02 Office action relating to structure-function relationships in proteins is simply not germane to whether one can make and use the polypeptide variants recited by the present claims [*i.e.*, Attwood et al. (Comput. Chem., 25(4):329-339, 2001)]. Likewise, the cited document relating to alleged difficulties in assigning protein function based on homology comparison is not relevant to making the claimed polynucleotide variants [*i.e.*, Ponting (Brief. Bioinform., 2(1):19-29, 2001)]. That is, regardless of the precise functional characteristics of the SEQ ID NO:1 and SEQ ID NO:2 variants, one can still make the claimed polynucleotide variants using the disclosure provided by the present Specification. The polynucleotides could then be used in, for example, diagnostic testing, drug discovery, expression profiling, etc., as discussed in the Bedilion Declaration, filed April 16, 2003.

Furthermore, the Board's attention is also directed to the enclosed reference by Brenner et al. ("Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships," Proc. Natl. Acad. Sci. USA (1998) 95:6073-6078). Through exhaustive analysis of a data set of proteins with known structural and functional relationships and with <90% overall sequence identity, Brenner et al. have determined that 30% identity is a reliable threshold for establishing evolutionary homology between two sequences aligned over at least 150 residues. (Brenner et al., pages 6073 and 6076.) Furthermore, local identity is particularly important in this case for assessing the significance of the alignments, as Brenner et al. further report that $\geq 40\%$ identity over at least 70 residues is reliable in signifying homology between proteins. (Brenner et al., page 6076.)

Claim 3 recites, *inter alia*, a polynucleotide encoding a polypeptide comprising “a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.” In accordance with Brenner et al, naturally occurring molecules may exist which could be characterized as MT-MDH proteins and which have as little as 30% identity over at least 150 residues to SEQ ID NO:1. The “95% variants” recited by the present claims have a variation that is far less than that of all potential MT-MDH proteins related to SEQ ID NO:1, i.e., those MT-MDH proteins having as little as 30% identity over at least 150 residues to SEQ ID NO:1. Therefore, one would expect the SEQ ID NO:1 variants recited by the present claims to have the functional activities of a MT-MDH protein.

As set forth in *In re Marzocchi*, 169 USPQ 367, 369 (CCPA 1971):

The first paragraph of § 112 ***requires nothing more than objective enablement.*** [emphasis added] How such a teaching is set forth, either by the use of illustrative examples or by broad terminology, is of no importance.

As a matter of Patent Office practice, then, a specification disclosure which contains a teaching of the manner and process of making and using the invention in terms which correspond in scope to those used in describing and defining the subject matter sought to be patented *must* be taken as in compliance with the enabling requirement of the first paragraph of § 112 *unless* there is reason to doubt the objective truth of the statements contained therein which must be relied on for enabling support.

Contrary to the standard set forth in *Marzocchi*, the Examiner has failed to provide any *reasons* why one would doubt that the guidance provided by the present Specification would enable one to make and use the recited variants of SEQ ID NO:1 or SEQ ID NO:2. Hence, a *prima facie* case for non-enablement has not been established with respect to the variants of SEQ ID NO:1 or SEQ ID NO:2.

For at least the above reasons, reversal of this rejection is requested.

(9) CONCLUSION

For at least the reasons set forth above, reversal of the rejections under 35 U.S.C. § 112, first paragraph, based on lack of an adequate written description and on lack of enablement, is respectfully requested.


If the USPTO determines that any additional fees are due, the Commissioner is hereby authorized to charge Deposit Account No. **09-0108**.

This brief is enclosed in triplicate.

Respectfully submitted,

INCYTE CORPORATION

Date: 06 January 2004




Richard C. Ekstrom

Reg. No. 37,027

Direct Dial Telephone: (650) 843-7352

Date: 1/06/04



Joel Harris

Reg. No. 44,743

Direct Dial Telephone: (650) 845-4866

3160 Porter Drive
Palo Alto, California 94304
Phone: (650) 855-0555
Fax: (650) 849-8886

Enclosures:

1. Brenner et al., Proc. Natl. Acad. Sci. U.S.A. 95:6073-78 (1998).

APPENDIX - CLAIMS ON APPEAL

3. An isolated polynucleotide encoding a polypeptide selected from the group consisting of:
- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1; and
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. A cell transformed with a recombinant polynucleotide of claim 6.
9. A method for producing a polypeptide encoded by a polynucleotide of claim 3, the method comprising:
- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide of claim 3, and
 - b) recovering the polypeptide so expressed.
12. An isolated polynucleotide selected from the group consisting of:
- a) a polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2,
 - b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2,
 - c) a polynucleotide having a sequence complementary to a polynucleotide of a),
 - d) a polynucleotide having a sequence complementary to a polynucleotide of b) and
 - e) an RNA equivalent of a)-d).



Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships

STEVEN E. BRENNER^{*†‡}, CYRUS CHOTHIA^{*}, AND TIM J. P. HUBBARD[§]

^{*}MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, United Kingdom; and [†]Sanger Centre, Wellcome Trust Genome Campus, Hinxton, Cambs CB10 1SA, United Kingdom

Communicated by David R. Davies, National Institute of Diabetes, Bethesda, MD, March 16, 1998 (received for review November 12, 1997)

ABSTRACT Pairwise sequence comparison methods have been assessed using proteins whose relationships are known reliably from their structures and functions, as described in the SCOP database [Murzin, A. G., Brenner, S. E., Hubbard, T. & Chothia C. (1995) *J. Mol. Biol.* 247, 536–540]. The evaluation tested the programs BLAST [Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403–410], WU-BLAST2 [Altschul, S. F. & Gish, W. (1996) *Methods Enzymol.* 266, 460–480], FASTA [Pearson, W. R. & Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. USA* 85, 2444–2448], and SSEARCH [Smith, T. F. & Waterman, M. S. (1981) *J. Mol. Biol.* 147, 195–197] and their scoring schemes. The error rate of all algorithms is greatly reduced by using statistical scores to evaluate matches rather than percentage identity or raw scores. The E-value statistical scores of SSEARCH and FASTA are reliable: the number of false positives found in our tests agrees well with the scores reported. However, the P-values reported by BLAST and WU-BLAST2 exaggerate significance by orders of magnitude. SSEARCH, FASTA $ktup = 1$, and WU-BLAST2 perform best, and they are capable of detecting almost all relationships between proteins whose sequence identities are $>30\%$. For more distantly related proteins, they do much less well; only one-half of the relationships between proteins with 20–30% identity are found. Because many homologs have low sequence similarity, most distant relationships cannot be detected by any pairwise comparison method; however, those which are identified may be used with confidence.

Sequence database searching plays a role in virtually every branch of molecular biology and is crucial for interpreting the sequences issuing forth from genome projects. Given the method's central role, it is surprising that overall and relative capabilities of different procedures are largely unknown. It is difficult to verify algorithms on sample data because this requires large data sets of proteins whose evolutionary relationships are known unambiguously and independently of the methods being evaluated. However, nearly all known homologs have been identified by sequence analysis (the method to be tested). Also, it is generally very difficult to know, in the absence of structural data, whether two proteins that lack clear sequence similarity are unrelated. This has meant that although previous evaluations have helped improve sequence comparison, they have suffered from insufficient, imperfectly characterized, or artificial test data. Assessment also has been problematic because high quality database sequence searching attempts to have both sensitivity (detection of homologs) and specificity (rejection of unrelated proteins); however, these complementary goals are linked such that increasing one causes the other to be reduced.

Sequence comparison methodologies have evolved rapidly, so no previously published tests have evaluated modern versions of programs commonly used. For example, parameters in BLAST (1) have changed, and WU-BLAST2 (2)—which produces gapped alignments—has become available. The latest version of FASTA (3) previously tested was 1.6, but the current release (version 3.0) provides fundamentally different results in the form of statistical scoring.

The previous reports also have left gaps in our knowledge. For example, there has been no published assessment of thresholds for scoring schemes more sophisticated than percentage identity. Thus, the widely discussed statistical scoring measures have never actually been evaluated on large databases of real proteins. Moreover, the different scoring schemes commonly in use have not been compared.

Beyond these issues, there is a more fundamental question: in an absolute sense, how well does pairwise sequence comparison work? That is, what fraction of homologous proteins can be detected using modern database searching methods?

In this work, we attempt to answer these questions and to overcome both of the fundamental difficulties that have hindered assessment of sequence comparison methodologies. First, we use the set of distant evolutionary relationships in the SCOP: Structural Classification of Proteins database (4), which is derived from structural and functional characteristics (5). The SCOP database provides a uniquely reliable set of homologs, which are known independently of sequence comparison. Second, we use an assessment method that jointly measures both sensitivity and specificity. This method allows straightforward comparison of different sequence searching procedures. Further, it can be used to aid interpretation of real database searches and thus provide optimal and reliable results.

Previous Assessments of Sequence Comparison. Several previous studies have examined the relative performance of different sequence comparison methods. The most encompassing analyses have been by Pearson (6, 7), who compared the three most commonly used programs. Of these, the Smith-Waterman algorithm (8) implemented in SSEARCH (3) is the oldest and slowest but the most rigorous. Modern heuristics have provided BLAST (1) the speed and convenience to make it the most popular program. Intermediate between these two is FASTA (3), which may be run in two modes offering either greater speed ($ktup = 2$) or greater effectiveness ($ktup = 1$). Pearson also considered different parameters for each of these programs.

To test the methods, Pearson selected two representative proteins from each of 67 protein superfamilies defined by the PIR database (9). Each was used as a query to search the database, and the matched proteins were marked as being homologous or unrelated according to their membership of PIR

Abbreviation: EPO, errors per query.

[†]Present address: Department of Structural Biology, Stanford University, Fairchild Building D-109, Stanford, CA 94305-5126

[‡]To whom reprint requests should be addressed. e-mail: brenner@hyper.stanford.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/956073-6\$2.00/0
PNAS is available online at <http://www.pnas.org>.

superfamilies. Pearson found that modern matrices and "ln-scaling" of raw scores improve results considerably. He also reported that the rigorous Smith-Waterman algorithm worked slightly better than FASTA, which was in turn more effective than BLAST.

Very large scale analyses of matrices have been performed (10), and Henikoff and Henikoff (11) also evaluated the effectiveness of BLAST and FASTA. Their test with BLAST considered the ability to detect homologs above a predetermined score but had no penalty for methods which also reported large numbers of spurious matches. The Henikoffs searched the SWISS-PROT database (12) and used PROSITE (13) to define homologous families. Their results showed that the BLOSUM62 matrix (14) performed markedly better than the extrapolated PAM-series matrices (15), which previously had been popular.

A crucial aspect of any assessment is the data that are used to test the ability of the program to find homologs. But in Pearson's and the Henikoffs' evaluations of sequence comparison, the correct results were effectively unknown. This is because the superfamilies in PIR and PROSITE are principally created by using the same sequence comparison methods which are being evaluated. Interdependency of data and methods creates a "chicken and egg" problem, and means for example, that new methods would be penalized for correctly identifying homologs missed by older programs. For instance, immunoglobulin variable and constant domains are clearly homologous, but PIR places them in different superfamilies. The problem is widespread: each superfamily in PIR 48.00 with a structural homolog is itself homologous to an average of 1.6 other PIR superfamilies (16).

To surmount these sorts of difficulties, Sander and Schneider (17) used protein structures to evaluate sequence comparison. Rather than comparing different sequence comparison algorithms, their work focused on determining a length-dependent threshold of percentage identity, above which all proteins would be of similar structure. A result of this analysis was the HSSP equation; it states that proteins with 25% identity over 80 residues will have similar structures, whereas shorter alignments require higher identity. (Other studies also have used structures (18-20), but these focused on a small number of model proteins and were principally oriented toward evaluating alignment accuracy rather than homology detection.)

A general solution to the problem of scoring comes from statistical measures (i.e., E-values and P-values) based on the extreme value distribution (21). Extreme value scoring was implemented analytically in the BLAST program using the Karlin and Altschul statistics (22, 23) and empirical approaches have been recently added to FASTA and SSEARCH. In addition to being heralded as a reliable means of recognizing significantly similar proteins (24, 25), the mathematical tractability of statistical scores "is a crucial feature of the BLAST algorithm" (1). The validity of this scoring procedure has been tested analytically and empirically (see ref. 2 and references in ref. 24). However, all large empirical tests used random sequences that may lack the subtle structure found within biological sequences (26, 27) and obviously do not contain any real homologs. Thus, although many researchers have suggested that statistical scores be used to rank matches (24, 25, 28), there have been no large rigorous experiments on biological data to determine the degree to which such rankings are superior.

A Database for Testing Homology Detection. Since the discovery that the structures of hemoglobin and myoglobin are very similar though their sequences are not (29), it has been apparent that comparing structures is a more powerful (if less convenient) way to recognize distant evolutionary relationships than comparing sequences. If two proteins show a high degree of similarity in their structural details and function, it

is very probable that they have an evolutionary relationship though their sequence similarity may be low.

The recent growth of protein structure information combined with the comprehensive evolutionary classification in the SCOP database (4, 5) have allowed us to overcome previous limitations. With these data, we can evaluate the performance of sequence comparison methods on real protein sequences whose relationships are known confidently. The SCOP database uses structural information to recognize distant homologs, the large majority of which can be determined unambiguously. These superfamilies, such as the globins or the immunoglobulins, would be recognized as related by the vast majority of the biological community despite the lack of high sequence similarity.

From SCOP, we extracted the sequences of domains of proteins in the Protein Data Bank (PDB) (30) and created two databases. One (PDB90D-B) has domains, which were all <90% identical to any other, whereas (PDB40D-B) had those <40% identical. The databases were created by first sorting all protein domains in SCOP by their quality and making a list. The highest quality domain was selected for inclusion in the database and removed from the list. Also removed from the list (and discarded) were all other domains above the threshold level of identity to the selected domain. This process was repeated until the list was empty. The PDB40D-B database contains 1,323 domains, which have 9,044 ordered pairs of distant relationships, or ~0.5% of the total 1,749,006 ordered pairs. In PDB90D-B, the 2,079 domains have 53,988 relationships, representing 1.2% of all pairs. Low complexity regions of sequence can achieve spurious high scores, so these were masked in both databases by processing with the SEG program (27) using recommended parameters: 12 1.8 2.0. The databases used in this paper are available from <http://sss.stanford.edu/sss/>, and databases derived from the current version of SCOP may be found at <http://scop.mrc-lmb.cam.ac.uk/scop/>.

Analyses from both databases were generally consistent, but PDB40D-B focuses on distantly related proteins and reduces the heavy overrepresentation in the PDB of a small number of families (31, 32), whereas PDB90D-B (with more sequences) improves evaluations of statistics. Except where noted otherwise, the distant homolog results here are from PDB40D-B. Although the precise numbers reported here are specific to the structural domain databases used, we expect the trends to be general.

Assessment Data and Procedure. Our assessment of sequence comparison may be divided into four different major categories of tests. First, using just a single sequence comparison algorithm at a time, we evaluated the effectiveness of different scoring schemes. Second, we assessed the reliability of scoring procedures, including an evaluation of the validity of statistical scoring. Third, we compared sequence comparison algorithms (using the optimal scoring scheme) to determine their relative performance. Fourth, we examined the distribution of homologs and considered the power of pairwise sequence comparison to recognize them. All of the analyses used the databases of structurally identified homologs and a new assessment criterion.

The analyses tested BLAST (1), version 1.4.9MP, and WU-BLAST2 (2), version 2.0a13MP. Also assessed was the FASTA package, version 3.0i76 (3), which provided FASTA and the SSEARCH implementation of Smith-Waterman (8). For SSEARCH and FASTA, we used BLOSUM45 with gap penalties -12/-1 (7, 16). The default parameters and matrix (BLOSUM62) were used for BLAST and WU-BLAST2.

The "Coverage Vs. Error" Plot. To test a particular protocol (comprising a program and scoring scheme), each sequence from the database was used as a query to search the database. This yielded ordered pairs of query and target sequences with associated scores, which were sorted, on the basis of their scores, from best to worst. The ideal method would have




FIG. 1. Coverage vs. error plots of different scoring schemes for SSEARCH Smith-Waterman. (A) Analysis of PDB40D-B database. (B) Analysis of PDB90D-B database. All of the proteins in the database were compared with each other using the SSEARCH program. The results of this single set of comparisons were considered using five different scoring schemes and assessed. The graphs show the coverage and errors per query (EPQ) for statistical scores, raw scores, and three measures using percentage identity. In the coverage vs. error plot, the x axis indicates the fraction of all homologs in the database (known from structure) which have been detected. Precisely, it is the number of detected pairs of proteins with the same fold divided by the total number of pairs from a common superfamily. PDB40D-B contains a total of 9,044 homologs, so a score of 10% indicates identification of 904 relationships. The y axis reports the number of EPQ. Because there are 1,323 queries made in the PDB40D-B all-vs.-all comparison, 13 errors corresponds to 0.01, or 1% EPQ. The y axis is presented on a log scale to show results over the widely varying degrees of accuracy which may be desired. The scores that correspond to the levels of EPQ and coverage are shown in Fig. 4 and Table 1. The graph demonstrates the trade-off between sensitivity and selectivity. As more homologs are found (moving to the right), more errors are made (moving up). The ideal method would be in the lower right corner of the graph, which corresponds to identifying many evolutionary relationships without selecting unrelated proteins. Three measures of percentage identity are plotted. Percentage identity within alignment is the degree of identity within the aligned region of the proteins, without consideration of the alignment length. Percentage identity within both is the number of identical residues in the aligned region as a percentage of the average length of the query and target proteins. The HSP equation (17) is $H = 290.15 / -0.542$ where l is length for $10 < l < 80$; $H > 100$ for $l < 10$; $H = 24.7$ for $l > 80$. The percentage identity HSP-adjusted score is the percent identity within the alignment minus H . Smith-Waterman raw scores and E-values were taken directly from the sequence comparison program.

perfect separation, with all of the homologs at the top of the list and unrelated proteins below. In practice, perfect separation is impossible to achieve so instead one is interested in drawing a threshold above which there are the largest number of related pairs of sequences consistent with an acceptable error rate.

Our procedure involved measuring the coverage and error for every threshold. Coverage was defined as the fraction of structurally determined homologs that have scores above the selected threshold; this reflects the sensitivity of a method. Errors per query (EPQ), an indicator of selectivity, is the number of nonhomologous pairs above the threshold divided by the number of queries. Graphs of these data, called coverage vs. error plots, were devised to understand how

protocols compare at different levels of accuracy. These graphs share effectively all of the beneficial features of Receiver Operating Characteristic (ROC) plots (33, 34) but better represent the high degrees of accuracy required in sequence comparison and the huge background of nonhomologs.

This assessment procedure is directly relevant to practical sequence database searching, for it provides precisely the information necessary to perform a reliable sequence database search. The EPQ measure places a premium on score consistency: that is, it requires scores to be comparable for different queries. Consistency is an aspect which has been largely




FIG. 2. Unrelated proteins with high percentage identity. Hemoglobin β -chain (PDB code 1hds chain b, ref. 38, Left) and cellulase E2 (PDB code 1tml, ref. 39, Right) have 39% identity over 64 residues, a level which is often believed to be indicative of homology. Despite this high degree of identity, their structures strongly suggest that these proteins are not related. Appropriately, neither the raw alignment score of 85 nor the E-value of 1.3 is significant. Proteins rendered by RASMOL (40).




FIG. 3. Length and percentage identity of alignments of unrelated proteins in PDB90D-B. Each pair of nonhomologous proteins found with SSEARCH is plotted as a point whose position indicates the length and the percentage identity within the alignment. Because alignment length and percentage identity are quantized, many pairs of proteins may have exactly the same alignment length and percentage identity. The line shows the HSP threshold (though it is intended to be applied with a different matrix and parameters).




FIG. 4. Reliability of statistical scores in PDB90D-B: Each line shows the relationship between reported statistical score and actual error rate for a different program. E-values are reported for SSEARCH and FASTA, whereas P-values are shown for BLAST and WU-BLAST. If the scoring were perfect, then the number of errors per query and the E-values would be the same, as indicated by the upper bold line. (P-values should be the same as EPO for small numbers, and diverges at higher values, as indicated by the lower bold line.) E-values from SSEARCH and FASTA are shown to have good agreement with EPO but underestimate the significance slightly. BLAST and WU-BLAST are overconfident, with the degree of exaggeration dependent upon the score. The results for PDB40D-B were similar to those for PDB90D-B despite the difference in number of homologs detected. This graph could be used to roughly calibrate the reliability of a given statistical score.

ignored in previous tests but is essential for the straightforward or automatic interpretation of sequence comparison results. Further, it provides a clear indication of the confidence that should be ascribed to each match. Indeed, the EPQ measure should approximate the expectation value reported by database searching programs, if the programs' estimates are accurate.

The Performance of Scoring Schemes. All of the programs tested could provide three fundamental types of scores. The first score is the percentage identity, which may be computed in several ways based on either the length of the alignment or the lengths of the sequences. The second is a "raw" or "Smith-Waterman" score, which is the measure optimized by the Smith-Waterman algorithm and is computed by summing the substitution matrix scores for each position in the alignment and subtracting gap penalties. In BLAST, a measure

related to this score is scaled into bits. Third is a statistical score based on the extreme value distribution. These results are summarized in Fig. 1.

Sequence Identity. Though it has been long established that percentage identity is a poor measure (35), there is a common rule-of-thumb stating that 30% identity signifies homology. Moreover, publications have indicated that 25% identity can be used as a threshold (17, 36). We find that these thresholds, originally derived years ago, are not supported by present results. As databases have grown, so have the possibilities for chance alignments with high identity; thus, the reported cutoffs lead to frequent errors. Fig. 2 shows one of the many pairs of proteins with very different structures that nonetheless have high levels of identity over considerable aligned regions. Despite the high identity, the raw and the statistical scores for such incorrect matches are typically not significant. The principal reasons percentage identity does so poorly seem to be that it ignores information about gaps and about the conservative or radical nature of residue substitutions.

From the PDB90D-B analysis in Fig. 3, we learn that 30% identity is a reliable threshold for this database only for sequence alignments of at least 150 residues. Because one unrelated pair of proteins has 43.5% identity over 62 residues, it is probably necessary for alignments to be at least 70 residues in length before 40% is a reasonable threshold, for a database of this particular size and composition.

At a given reliability, scores based on percentage identity detect just a fraction of the distant homologs found by statistical scoring. If one measures the percentage identity in the aligned regions without consideration of alignment length, then a negligible number of distant homologs are detected. Use of the HSSP equation improves the value of percentage identity, but even this measure can find only 4% of all known homologs at 1% EPQ. In short, percentage identity discards most of the information measured in a sequence comparison.

Raw Scores. Smith-Waterman raw scores perform better than percentage identity (Fig. 1), but ln-scaling (7) provided no notable benefit in our analysis. It is necessary to be very precise when using either raw or bit scores because a 20% change in cutoff score could yield a tenfold difference in EPQ. However, it is difficult to choose appropriate thresholds because the reliability of a bit score depends on the lengths of the proteins matched and the size of the database. Raw score thresholds also are affected by matrix and gap parameters.

Statistical Scores. Statistical scores were introduced partly to overcome the problems that arise from raw scores. This scoring scheme provides the best discrimination between homologous proteins and those which are unrelated. Most





FIG. 5. Coverage vs. error plots of different sequence comparison methods: Five different sequence comparison methods are evaluated, each using statistical scores (E- or P-values). (A) PDB40D-B database. In this analysis, the best method is the slow SSEARCH, which finds 18% of relationships at 1% EPQ. FASTA ktup = 1 and WU-BLAST2 are almost as good. (B) PDB90D-B database. The quick WU-BLAST2 program provides the best coverage at 1% EPQ on this database, although at higher levels of error it becomes slightly worse than FASTA ktup = 1 and SSEARCH.

likely, its power can be attributed to its incorporation of more information than any other measure; it takes account of the full substitution and gap data (like raw scores) but also has details about the sequence lengths and composition and is scaled appropriately.

We find that statistical scores are not only powerful, but also easy to interpret. SSEARCH and FASTA show close agreement between statistical scores and actual number of errors per query (Fig. 4). The expectation value score gives a good, slightly conservative estimate of the chances of the two sequences being found at random in a given query. Thus, an E-value of 0.01 indicates that roughly one pair of nonhomologs of this similarity should be found in every 100 different queries. Neither raw scores nor percentage identity can be interpreted in this way, and these results validate the suitability of the extreme value distribution for describing the scores from a database search.

The P-values from BLAST also should be directly interpretable but were found to overstate significance by more than two orders of magnitude for 1% EPO for this database. Nonetheless, these results strongly suggest that the analytic theory is fundamentally appropriate. WU-BLAST2 scores were more reliable than those from BLAST, but also exaggerate expected confidence by more than an order of magnitude at 1% EPO.

Overall Detection of Homologs and Comparison of Algorithms. The results in Fig. 5A and Table 1 show that pairwise sequence comparison is capable of identifying only a small fraction of the homologous pairs of sequences in PDB40D-B. Even SSEARCH with E-values, the best protocol tested, could find only 18% of all relationships at a 1% EPO. BLAST, which identifies 15%, was the worst performer, whereas FASTA ktup = 1 is nearly as effective as SSEARCH. FASTA ktup = 2 and WU-BLAST2 are intermediate in their ability to detect homologs. Comparison of different algorithms indicates that those capable of identifying more homologs are generally slower. SSEARCH is 25 times slower than BLAST and 6.5 times slower than FASTA ktup = 1. WU-BLAST2 is slightly faster than FASTA ktup = 2, but the latter has more interpretable scores.

In PDB90D-B, where there are many close relationships, the best method can identify only 38% of structurally known homologs (Fig. 5B). The method which finds that many relationships is WU-BLAST2. Consequently, we infer that the differences between FASTA ktup = 1, SSEARCH, and WU-BLAST2 programs are unlikely to be significant when compared with variation in database composition and scoring reliability.

Fig. 6 helps to explain why most distant homologs cannot be found by sequence comparison: a great many such relationships have no more sequence identity than would be expected by chance. SSEARCH with E-values can recognize >90% of the homologous pairs with 30–40% identity. In this region, there are 30 pairs of homologous proteins that do not have significant E-values, but 26 of these involve sequences with <50 residues. Of sequences having 25–30% identity, 75% are identified by SSEARCH E-values. However, although the number of homologs grows at lower levels of identity, the detection falls off sharply: only 40% of homologs with 20–25% identity




FIG. 6. Distribution and detection of homologs in PDB40D-B. Bars show the distribution of homologous pairs PDB40D-B according to their identity (using the measure of identity in both). Filled regions indicate the number of these pairs found by the best database searching method (SSEARCH with E-values) at 1% EPO. The PDB40D-B database contains proteins with <40% identity, and as shown on this graph, most structurally identified homologs in the database have diverged extremely far in sequence and have <20% identity. Note that the alignments may be inaccurate, especially at low levels of identity. Filled regions show that SSEARCH can identify most relationships that have 25% or more identity, but its detection wanes sharply below 25%. Consequently, the great sequence divergence of most structurally identified evolutionary relationships effectively defeats the ability of pairwise sequence comparison to detect them.

are detected and only 10% of those with 15–20% can be found. These results show that statistical scores can find related proteins whose identity is remarkably low; however, the power of the method is restricted by the great divergence of many protein sequences.

After completion of this work, a new version of pairwise BLAST was released: BLASTGP (37). It supports gapped alignments, like WU-BLAST2, and dispenses with sum statistics. Our initial tests on BLASTGP using default parameters show that its E-values are reliable and that its overall detection of homologs was substantially better than that of ungapped BLAST, but not quite equal to that of WU-BLAST2.

CONCLUSION

The general consensus amongst experts (see refs. 7, 24, 25, 27 and references therein) suggests that the most effective sequence searches are made by (i) using a large current database in which the protein sequences have been complexity masked and (ii) using statistical scores to interpret the results. Our experiments fully support this view.

Our results also suggest two further points. First, the E-values reported by FASTA and SSEARCH give fairly accurate estimates of the significance of each match, but the P-values provided by BLAST and WU-BLAST2 underestimate the true

Table 1. Summary of sequence comparison methods with PDB40D-B

Method	Relative Time*	1% EPO Cutoff	Coverage at 1% EPO
SSEARCH % identity: within alignment	25.5	>70%	<0.1
SSEARCH % identity: within both	25.5	34%	3.0
SSEARCH % identity: HSSP-scaled	25.5	35% (HSSP = 9.8)	4.0
SSEARCH Smith-Waterman raw scores	25.5	14%	10.5
SSEARCH E-values	25.5	0.03	18.4
FASTA ktup = 1 E-values	3.9	0.03	17.9
FASTA ktup = 2 E-values	1.4	0.03	16.7
WU-BLAST2 P-values	1.1	0.003	17.5
BLAST P-values	1.0	0.00016	14.8

*Times are from large database searches with genome proteins.

extent of errors. Second, SSEARCH, WU-BLAST2, and FASTA ktup = 1 perform best, though BLAST and FASTA ktup = 2 detect most of the relationships found by the best procedures and are appropriate for rapid initial searches.

The homologous proteins that are found by sequence comparison can be distinguished with high reliability from the huge number of unrelated pairs. However, even the best database searching procedures tested fail to find the large majority of distant evolutionary relationships at an acceptable error rate. Thus, if the procedures assessed here fail to find a reliable match, it does not imply that the sequence is unique; rather, it indicates that any relatives it might have are distant ones.**

**Additional and updated information about this work, including supplementary figures, may be found at <http://sss.stanford.edu/sss/>.

The authors are grateful to Drs. A. G. Murzin, M. Levitt, S. R. Eddy, and G. Mitchison for valuable discussion. S.E.B. was principally supported by a St. John's College (Cambridge, UK) Benefactors' Scholarship and by the American Friends of Cambridge University. S.E.B. dedicates his contribution to the memory of Rabbi Albert T. and Clara S. Bilgray.

1. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403-410.
2. Altschul, S. F. & Gish, W. (1996) *Methods Enzymol.* 266, 460-480.
3. Pearson, W. R. & Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. USA* 85, 2444-2448.
4. Murzin, A. G., Brenner, S. E., Hubbard, T., & Chothia, C. (1995) *J. Mol. Biol.* 247, 536-540.
5. Brenner, S. E., Chothia, C., Hubbard, T. J. P., & Murzin, A. G. (1996) *Methods Enzymol.* 266, 635-643.
6. Pearson, W. R. (1991) *Genomics* 11, 635-650.
7. Pearson, W. R. (1995) *Protein Sci.* 4, 1145-1160.
8. Smith, T. F. & Waterman, M. S. (1981) *J. Mol. Biol.* 147, 195-197.
9. George, D. G., Hunt, L. T., & Barker, W. C. (1996) *Methods Enzymol.* 266, 41-59.
10. Vogt, G., Etzold, T., & Argos, P. (1995) *J. Mol. Biol.* 249, 816-831.
11. Henikoff, S. & Henikoff, J. G. (1993) *Proteins* 17, 49-61.
12. Bairoch, A. & Apweiler, R. (1996) *Nucleic Acids Res.* 24, 21-25.
13. Bairoch, A., Bucher, P., & Hofmann, K. (1996) *Nucleic Acids Res.* 24, 189-196.
14. Henikoff, S. & Henikoff, J. G. (1992) *Proc. Natl. Acad. Sci. USA* 89, 10915-10919.
15. Dayhoff, M., Schwartz, R. M., & Orcutt, B. C. (1978) in *Atlas of Protein Sequence and Structure*, ed. Dayhoff, M. (National Bio-
medical Research Foundation, Silver Spring, MD), Vol. 5, Suppl. 3, pp. 345-352.
16. Brenner, S. E. (1996) Ph.D. thesis. (University of Cambridge, UK).
17. Sander, C. & Schneider, R. (1991) *Proteins* 9, 56-68.
18. Johnson, M. S. & Overington, J. P. (1993) *J. Mol. Biol.* 233, 716-738.
19. Barton, G. J. & Sternberg, M. J. E. (1987) *Protein Eng.* 1, 89-94.
20. Lesk, A. M., Levitt, M., & Chothia, C. (1986) *Protein Eng.* 1, 77-78.
21. Arratia, R., Gordon, L., & M. W. (1986) *Ann. Stat.* 14, 971-993.
22. Karlin, S. & Altschul, S. F. (1990) *Proc. Natl. Acad. Sci. USA* 87, 2264-2268.
23. Karlin, S. & Altschul, S. F. (1993) *Proc. Natl. Acad. Sci. USA* 90, 5873-5877.
24. Altschul, S. F., Boguski, M. S., Gish, W., & Wootton, J. C. (1994) *Nat. Genet.* 6, 119-129.
25. Pearson, W. R. (1996) *Methods Enzymol.* 266, 227-258.
26. Lipman, D. J., Wilbur, W. J., Smith, T. F., & Waterman, M. S. (1984) *Nucleic Acids Res.* 12, 215-226.
27. Wootton, J. C. & Federhen, S. (1996) *Methods Enzymol.* 266, 554-571.
28. Waterman, M. S. & Vingron, M. (1994) *Stat. Science* 9, 367-381.
29. Perutz, M. F., Kendrew, J. C., & Watson, H. C. (1965) *J. Mol. Biol.* 13, 669-678.
30. Abola, E. E., Bernstein, F. C., Bryant, S. H., Koetzle, T. F., & Weng, J. (1987) in *Crystallographic Databases: Information Content, Software Systems, Scientific Applications*, eds. Allen, F. H., Bergerhoff, G., & Sievers, R. (Data Comm. Intl. Union Crystallogr., Cambridge, UK), pp. 107-132.
31. Brenner, S. E., Chothia, C., & Hubbard, T. J. P. (1997) *Curr. Opin. Struct. Biol.* 7, 369-376.
32. Orengo, C., Michie, A., Jones, S., Jones, D. T., Swindells, M. B., & Thornton, J. (1997) *Structure (London)* 5, 1093-1108.
33. Zweig, M. H. & Campbell, G. (1993) *Clin. Chem.* 39, 561-577.
34. Gribskov, M. & Robinson, N. L. (1996) *Comput. Chem.* 20, 25-33.
35. Fitch, W. M. (1966) *J. Mol. Biol.* 16, 9-16.
36. Chung, S. Y. & Subbiah, S. (1996) *Structure (London)* 4, 1123-1127.
37. Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997) *Nucleic Acids Res.* 25, 3389-3402.
38. Girling, R., Schmidt, W., Jr., Houston, T., Amma, E., & Huisman, T. (1979) *J. Mol. Biol.* 131, 417-433.
39. Specio, M., Wilson, D., & Karplus, P. (1993) *Biochemistry* 32, 9906-9916.
40. Savle, R. A. & Milner-White, E. J. (1995) *Trends Biochem. Sci.* 20, 374-376.



Docket No.: PF-0379-1 DIV

Response Under 37 C.F.R. 1.116 - Expedited Procedure
Examining Group 1652

Certificate of Mailing

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Mail Stop Appeal Brief Patents, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450 on _____

By: [Signature] Printed: _____

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
BEFORE THE BOARD OF PATENT APPEALS AND INTERFERENCES**

In re Application of: Bandman et al.

Title: HUMAN MITOCHONDRIAL MALATE DEHYDROGENASE

Serial No.: 09/915,694

Filing Date: July 25, 2001

Examiner: Fronda, C.

Group Art Unit: 1652

Mail Stop Appeal Brief-Patents
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

BRIEF ON APPEAL

Sir:

Further to the Notice of Appeal filed on September 30, 2003, and received by the USPTO on October 6, 2003, herewith are three copies of Appellants' Brief on Appeal. Appellants hereby request a one month extension of time in order to file this Brief. Authorized fees include the statutory fee of \$110 for a one-month extension of time, as well as the \$ 330.00 fee for the filing of this Brief.

This is an appeal from the decision of the Examiner finally rejecting claims 3, 6, 7, 9 and 12 of the above-identified application.

(1) REAL PARTY IN INTEREST

The above-identified application is assigned of record to Incyte Pharmaceuticals, Inc. (now Incyte Corporation, formerly known as Incyte Genomics, Inc.) (Reel 9774, Frame 0975), which is the real party in interest herein.

(2) RELATED APPEALS AND INTERFERENCES

Appellants, their legal representative and the assignee are not aware of any related appeals or interferences which will directly affect or be directly affected by or have a bearing on the Board's decision in the instant appeal.

(3) STATUS OF THE CLAIMS

Claims rejected: Claims 3, 6, 7, 9 and 12
Claims allowed: (none)
Claims objected to: Claims 4, 5 and 10.
Claims canceled: Claims 1, 2, 8, 11, 13, 17-27, 30-45.
Claims withdrawn: Claims 14-16, 28 and 29
Claims on Appeal: Claims 3, 6, 7, 9 and 12 (A copy of the claims on appeal, as amended, can be found in the attached Appendix).

(4) STATUS OF AMENDMENTS AFTER FINAL

There were no amendments made after final.

(5) SUMMARY OF THE INVENTION

Embodiments of the present invention are directed, *inter alia*, to polynucleotides encoding mitochondrial malate dehydrogenase (MT-MDH). In particular embodiments, the mitochondrial malate dehydrogenases are selected from among amino acid sequences comprising SEQ ID NO:1. These polypeptides have strong chemical and structural homology with known mitochondrial malate dehydrogenases. For example:

MT-MDH is 338 amino acids in length and has two potential N-glycosylation sites at residues N-117 and N-145, seven potential casein kinase II phosphorylation sites at T-54, S-69, T-109, T-170, S-261, S-309, and S-310, four potential protein kinase C phosphorylation sites at residues T-213, T-227, S-326, and T-336, a mitochondrial malate dehydrogenase active site signature between residues V-169 and V-181, and a transit peptide sequence from residues M-1 to N-24. As shown in Figures 2A and 2B, MT-MDH has chemical and structural homology with murine mitochondrial malate dehydrogenase (GI 56643; SEQ ID NO:3) and porcine mitochondrial malate dehydrogenase (GI 164541; SEQ ID NO:4). In particular, MT-MDH and murine

mitochondrial mitochondrial malate dehydrogenase share 94% identity, share both potential N-glycosylation sites, six potential casein kinase II sites, three potential protein kinase C sites, the mitochondrial malate dehydrogenase active site signature, and the transit peptide sequence. As illustrated by Figures 3A and 3B, respectively, MT-MDH and murine mitochondrial mitochondrial malate dehydrogenase (SEQ ID NO:3) have similar isoelectric points (pI = 8.8). As illustrated by Figures 4A and 4B, MT-MDH contains potential NAD(H) and NADP(H) binding site motifs. Northern analysis shows the expression of this sequence in various libraries, at least 49% of which are immortalized or cancerous and at least 24% of which involve immune response. Of particular note is the expression of MT-MDH in fetal tissues; in cardiovascular, gut, nervous, and reproductive tissues; and in secretory and hematopoietic tissues. (Specification at page 14, line 29 to page 15, line 17).

The polynucleotides of the present invention have a variety of utilities. For example, they can be used for the diagnosis, prevention, or treatment of vesicle trafficking, immunological and neoplastic disorders. (See the Specification e.g., at page 14, lines 18-21)

(6) ISSUES

1. Whether claims 3, 6, 7, 9 and 12 meet the written description requirement of 35 U.S.C. § 112, first paragraph.
2. Whether claims 3, 6, 7, 9 and 12 meet the enablement requirement of 35 U.S.C. § 112, first paragraph.

(7) GROUPING OF THE CLAIMS

As to Issue 1

All of the claims on appeal are grouped together.

As to Issue 2

All of the claims on appeal are grouped together.

(8) APPELLANTS' ARGUMENTS

Issue 1- Written description rejection under 35 U.S.C. § 112, first paragraph

Claims 3, 6, 7, 9 and 12 were rejected under 35 U.S.C. §112, first paragraph, as allegedly

“containing subject matter which was not described in the specification in such a way as to reasonably convey to one skilled in the relevant art the inventor(s) at the time the application was filed, had possession of the claimed invention.” (06/20/03 Office Action, at page 4) In making the rejection, the Examiner asserts that:

There is no disclosure of any particular structure to function/activity relationship in the single disclosed species. The specification also fails to described additional representative species of these polynucleotides by any identifying structural characteristics or properties for which no predictability of structure is apparent. Given this lack of additional representative species as encompassed by the claims, Applicants have failed to sufficiently disclose the invention in such full, clear, concise and exact terms that a skilled artisan would recognize Applicants were in possession of the claimed invention. (12/17/02 Office Action, at pages 4-5).

This rejection is improper, as the claims define subject matter which is described in the Specification in such a way as to reasonably convey to one skilled in the art that the inventors had possession of the claimed subject matter at the time the application was filed. The requirements necessary to fulfill the written description requirement of 35 U.S.C. § 112, first paragraph, are well established by case law:

... the applicant must also convey with reasonable clarity to those skilled in the art that, as of the filing date sought, he or she was in possession of *the invention*. The invention is, for purposes of the “written description” inquiry, *whatever is now claimed*. *Vas-Cath, Inc. v. Mahurkar*, 19 U.S.P.Q.2d 1111, 1117 (Fed. Cir. 1991).

The Board’s attention is also drawn to the Patent and Trademark Office’s own Examination of Patent Applications Under the 35 U.S.C. § 112, para. 1”, published January 5, 2001, which provide that:

An applicant may also show that an invention is complete by disclosure of sufficiently detailed, relevant identifying characteristics⁴² which provide evidence that applicant was in possession of the claimed invention,⁴³ i.e., complete or partial structure, other physical and/or chemical properties, functional characteristics when coupled with a known or disclosed correlation between function and structure, or some combination of such characteristics.⁴⁴ What is conventional or well known to one of ordinary skill in the art need not be disclosed in detail.⁴⁵ If a skilled artisan would have understood the inventor to be in possession of the claimed invention at the time of filing, even if every nuance of

the claims is not explicitly described in the specification, then the adequate description requirement is met.⁴⁶

Thus, the written description standard is fulfilled by both what is specifically disclosed and what is conventional or well known to one skilled in the art.

A. The specification provides an adequate written description of the claimed "variants" of SEQ ID NO:1 and SEQ ID NO:2.

The subject matter encompassed by claims 3, 6, 7, 9 and 12 is either disclosed by the specification or is conventional or well known to one skilled in the art.

First note that the "variant" language of independent claim 3 recites a polynucleotide encoding "a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1" and the "variant" language of independent claim 12 recites "a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2."

The amino acid sequence of SEQ ID NO:1 and the polynucleotide sequence of SEQ ID NO:2 are explicitly disclosed in the specification. See, for example, the Sequence Listing. Variants of SEQ ID NO:1 and SEQ ID NO:2 are described in the Specification at, for example, page 4, lines 13-14; page 7, lines 4-7 and 12-18; and page 15, lines 18-27.

One of ordinary skill in the art would recognize polynucleotide sequences which are variants having a polynucleotide sequence at least 95% to SEQ ID NO:2, or which encode polypeptide variants having an amino acid sequence at least 95% identical to SEQ ID NO:1. Given any naturally occurring polynucleotide sequence, it would be routine for one of skill in the art to recognize whether it was a variant of SEQ ID NO:2, or whether it encoded a variant of SEQ ID NO:1. Accordingly, the specification provides an adequate written description of the recited polynucleotide variants of SEQ ID NO:2 and polynucleotides encoding polypeptide variants of SEQ ID NO:1.

1. The present claims specifically define the claimed genus through the recitation of chemical structure

Court cases in which "DNA claims" have been at issue commonly emphasize that the

recitation of structural features or chemical or physical properties are important factors to consider in a written description analysis of such claims. For example, in *Fiers v. Revel*, 25 U.S.P.Q.2d 1601, 1606 (Fed. Cir. 1993), the court stated that:

If a conception of a DNA requires a precise definition, such as by structure, formula, chemical name or physical properties, as we have held, then a description also requires that degree of specificity.

In a number of instances in which claims to DNA have been found invalid, the courts have noted that the claims attempted to define the claimed DNA in terms of functional characteristics without any reference to structural features. As set forth by the court in *University of California v. Eli Lilly and Co.*, 43 U.S.P.Q.2d 1398, 1406 (Fed. Cir. 1997):

In claims to genetic material, however, a generic statement such as "vertebrate insulin cDNA" or "mammalian insulin cDNA," without more, is not an adequate written description of the genus because it does not distinguish the claimed genus from others, except by function.

Thus, the mere recitation of functional characteristics of a DNA, without the definition of structural features, has been a common basis by which courts have found invalid claims to DNA. For example, in *Lilly*, 43 U.S.P.Q.2d at 1407, the court found invalid for violation of the written description requirement the following claim of U.S. Patent No. 4,652,525:

1. A recombinant plasmid replicable in procaryotic host containing within its nucleotide sequence a subsequence having the structure of the reverse transcript of an mRNA of a vertebrate, which mRNA encodes insulin.

In *Fiers*, 25 U.S.P.Q.2d at 1603, the parties were in an interference involving the following count:

A DNA which consists essentially of a DNA which codes for a human fibroblast interferon-beta polypeptide.

Party Revel in the *Fiers* case argued that its foreign priority application contained an adequate written description of the DNA of the count because that application mentioned a potential method for isolating the DNA. The Revel priority application, however, did not have a description of any particular DNA structure corresponding to the DNA of the count. The court therefore found that the Revel priority application lacked an adequate written description of the

subject matter of the count.

Thus, in *Lilly* and *Fiers*, nucleic acids were defined on the basis of functional characteristics and were found not to comply with the written description requirement of 35 U.S.C. §112; *i.e.*, "an mRNA of a vertebrate, which mRNA encodes insulin" in *Lilly*, and "DNA which codes for a human fibroblast interferon-beta polypeptide" in *Fiers*. In contrast to the situation in *Lilly* and *Fiers*, the claims at issue in the present application define polynucleotides and polypeptides in terms of chemical structure, rather than on functional characteristics. For example, the "variant language" of independent claims 3 and 12 recite chemical structure to define the claimed genus:

3. An isolated polynucleotide encoding a polypeptide selected from the group consisting of:.....
b) a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.

12. An isolated polynucleotide selected from the group consisting of:
b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2,....

From the above it should be apparent that the claims of the subject application are fundamentally different from those found invalid in *Lilly* and *Fiers*. The subject matter of the present claims is defined in terms of the chemical structure of SEQ ID NO:1 and SEQ ID NO:2. In the present case, there is no reliance merely on a description of functional characteristics of the polynucleotides and polypeptides recited by the claims. In fact, there is no recitation of functional characteristics. Moreover, if such functional recitations were included, it would add to the structural characterization of the recited polynucleotides. The polynucleotides defined in the claims of the present application recite structural features, and cases such as *Lilly* and *Fiers* stress that the recitation of structure is an important factor to consider in a written description analysis of claims of this type. By failing to base its written description inquiry "on whatever is now claimed," the Office Action fails to provide an appropriate analysis of the present claims and how they differ from those found not to satisfy the written description requirement in *Lilly* and *Fiers*.

2. The present claims do not define a genus which is "highly variant"

Furthermore, the claims at issue do not describe a genus which could be characterized as "highly variant." Available evidence illustrates that the claimed genus is of narrow scope.

In support of this assertion, the Board's attention is directed to the enclosed reference by Brenner et al. ("Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships," Proc. Natl. Acad. Sci. USA (1998) 95:6073-6078). Through exhaustive analysis of a data set of proteins with known structural and functional relationships and with <90% overall sequence identity, Brenner et al. have determined that 30% identity is a reliable threshold for establishing evolutionary homology between two sequences aligned over at least 150 residues. (Brenner et al., pages 6073 and 6076.) Furthermore, local identity is particularly important in this case for assessing the significance of the alignments, as Brenner et al. further report that $\geq 40\%$ identity over at least 70 residues is reliable in signifying homology between proteins. (Brenner et al., page 6076.)

The present application is directed, *inter alia*, to polynucleotides encoding mitochondrial malate dehydrogenases, including polynucleotides encoding mitochondrial malate dehydrogenases related to the amino acid sequence of SEQ ID NO:1. In accordance with Brenner et al, naturally occurring molecules may exist which could be characterized as mitochondrial malate dehydrogenases and which have as little as 30% identity over at least 150 residues to SEQ ID NO:1. The "variant language" of the present claims recites, for example, polynucleotides encoding a polypeptide comprising "a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1" (note that SEQ ID NO:1 has 338 amino acid residues). This variation is far less than that of polynucleotides encoding all potential mitochondrial malate dehydrogenases related to SEQ ID NO:1, i.e., those mitochondrial malate dehydrogenases having as little as 30% identity over at least 150 residues to SEQ ID NO:1.

3. The state of the art at the time of the present invention is further advanced than at the time of the *Lilly* and *Fiers* applications

In the *Lilly* case, claims of U.S. Patent No. 4,652,525 were found invalid for failing to comply with the written description requirement of 35 U.S.C. §112. The '525 patent claimed the

benefit of priority of two applications, Application Serial No. 801,343 filed May 27, 1977, and Application Serial No. 805,023 filed June 9, 1977. In the *Fiers* case, party Revel claimed the benefit of priority of an Israeli application filed on November 21, 1979. Thus, the written description inquiry in those case was based on the state of the art at essentially at the "dark ages" of recombinant DNA technology.

The present application has a priority date of September 3, 1997. Much has happened in the development of recombinant DNA technology in the 18 or more years from the time of filing of the applications involved in *Lilly* and *Fiers* and the present application. For example, the technique of polymerase chain reaction (PCR) was invented. Highly efficient cloning and DNA sequencing technology has been developed. Large databases of protein and nucleotide sequences have been compiled. Much of the raw material of the human and other genomes has been sequenced. With these remarkable advances one of skill in the art would recognize that, given the sequence information of SEQ ID NO:1 and SEQ ID NO:2, and the additional extensive detail provided by the subject application, the present inventors were in possession of the claimed polynucleotide variants at the time of filing of this application.

4. Summary

The 12/17/02 and 6/30/03 Office Actions fail to base the written description inquiry "on whatever is now claimed." Consequently, the Office Actions do not provide an appropriate analysis of the present claims and how they differ from those found not to satisfy the written description requirement in cases such as *Lilly* and *Fiers*. In particular, the claims of the subject application are fundamentally different from those found invalid in *Lilly* and *Fiers*. The subject matter of the present claims is defined in terms of the chemical structure of SEQ ID NO:1 and SEQ ID NO:2. The courts have stressed that structural features are important factors to consider in a written description analysis of claims to nucleic acids and proteins. In addition, the genus of polynucleotides defined by the present claims is adequately described, as evidenced by Brenner et al. Furthermore, there have been remarkable advances in the state of the art since the *Lilly* and *Fiers* cases, and these advances were given no consideration whatsoever in the position set forth by the Office Action.

For at least the reasons set forth above, the specification provides an adequate written description of the claimed subject matter, and this rejection should be reversed.

Issue 2 – Enablement rejection under 35 U.S.C. § 112, first paragraph

Claims 3, 6, 7, 9 and 12 stand rejected under 35 U.S.C. 112, first paragraph allegedly for lacking an enabling disclosure with respect to variants of SEQ ID NO:1. The Examiner has specifically stated that these claims contain “subject matter which was not described in the specification in such a way as to enable one skilled in the art to which it pertains, or with which it is most nearly connected, to make and/or use the invention.” (06/30/03 Office Action, at page 2-3)

In making the rejection, the Examiner asserts that:

“The specification does not provide guidance with respect to the specific structural/catalytic amino acids and the structural motifs essential for enzyme structure and activity function which cannot be altered. Thus searching for the specific nucleotides to change (deletion, insertion, substitution, or combinations thereof) in a polynucleotide is well outside the realm of routine experimentation and predictability in the art” (12/17/02 Office Action, at page 6).

The Examiner makes a similar assertion in the 06/30/03 Office Action that “[t]he specification does not teach the specific amino acids that can be altered and yet still retain enzyme activity.” (6/30/03 Office Action, at page 3).

The first paragraph of 35 U.S.C. §112 requires that the Specification describe how to make and use the claimed subject matter. That requirement has been met in the present application. In particular, the Specification describes how to make and use naturally-occurring polypeptide variants of SEQ ID NO:1 and polynucleotides encoding such variants.

Independent claim 3 recites not only that the “variant” polynucleotides encode polypeptides that are at least 95% identical to SEQ ID NO:1, but also have “*a naturally-occurring amino acid sequence.*” Through the process of natural selection, nature will have determined the appropriate amino acid sequences. Given the information provided by SEQ ID NO:1 (the amino acid sequence of MT-MDH) and SEQ ID NO:2 (the polynucleotide sequence encoding MT-MDH), one of skill in the art would be able to routinely obtain a polynucleotide encoding a polypeptide comprising “a naturally-occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.” Likewise for the “variant” polynucleotides defined by independent claim 12: “a polynucleotide comprising a naturally

occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2." For example, the identification of relevant polynucleotides could be performed by hybridization and/or PCR techniques that were well-known to those skilled in the art at the time the subject application was filed and/or described throughout the Specification of the instant application. For example:

The terms "stringent conditions" or "stringency", as used herein, refer to the conditions for hybridization as defined by the nucleic acid, salt, and temperature. These conditions are well known in the art and may be altered in order to identify or detect identical or related polynucleotide sequences. Numerous equivalent conditions comprising either low or high stringency depend on factors such as the length and nature of the sequence (DNA, RNA, base composition), nature of the target (DNA, RNA, base composition), milieu (in solution or immobilized on a solid substrate), concentration of salts and other components (e.g., formamide, dextran sulfate and/or polyethylene glycol), and temperature of the reactions (within a range from about 5°C below the melting temperature of the probe to about 20°C to 25°C below the melting temperature). One or more factors may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions. (Specification at page 13, lines 11-21)

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding MT-MDH or closely related molecules, may be used to identify nucleic acid sequences which encode MT-MDH. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding MT-MDH, alleles, or related sequences. (Specification at page 37, line 25 to page 38, line 3)

Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the MT-MDH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring MT-MDH. (Specification at page 38, lines 4-8)

See also Example VI, at page 50.

Thus, one skilled in the art need not make and test vast numbers of polypeptides that are based on the amino acid sequence of SEQ ID NO:1. Instead, one skilled in the art need only screen a cDNA library or use appropriate PCR conditions to identify relevant polynucleotides/polypeptides that already exist in nature. By adjusting the nature of the probe or nucleic acid (*i.e.*, non-conserved, conserved or highly conserved) and the conditions of hybridization (maximum, high, intermediate or low stringency), one can obtain variant polynucleotides of SEQ ID NO:2 which, in turn, will allow one to make the variant polypeptides of SEQ ID NO:1 recited by the present claims. Furthermore, the Specification sets forth an assay for measuring malate dehydrogenase activity (Example X at page 52, lines 11-18).

Accordingly, the document cited by the Examiner in the 12/17/02 Office action relating to structure-function relationships in proteins is simply not germane to whether one can make and use the polypeptide variants recited by the present claims [*i.e.*, Attwood et al. (Comput. Chem., 25(4):329-339, 2001)]. Likewise, the cited document relating to alleged difficulties in assigning protein function based on homology comparison is not relevant to making the claimed polynucleotide variants [*i.e.*, Ponting (Brief. Bioinform., 2(1):19-29, 2001)]. That is, regardless of the precise functional characteristics of the SEQ ID NO:1 and SEQ ID NO:2 variants, one can still make the claimed polynucleotide variants using the disclosure provided by the present Specification. The polynucleotides could then be used in, for example, diagnostic testing, drug discovery, expression profiling, etc., as discussed in the Bedilion Declaration, filed April 16, 2003.

Furthermore, the Board's attention is also directed to the enclosed reference by Brenner et al. ("Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships," Proc. Natl. Acad. Sci. USA (1998) 95:6073-6078). Through exhaustive analysis of a data set of proteins with known structural and functional relationships and with <90% overall sequence identity, Brenner et al. have determined that 30% identity is a reliable threshold for establishing evolutionary homology between two sequences aligned over at least 150 residues. (Brenner et al., pages 6073 and 6076.) Furthermore, local identity is particularly important in this case for assessing the significance of the alignments, as Brenner et al. further report that $\geq 40\%$ identity over at least 70 residues is reliable in signifying homology between proteins. (Brenner et al., page 6076.)

Claim 3 recites, *inter alia*, a polynucleotide encoding a polypeptide comprising “a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.” In accordance with Brenner et al, naturally occurring molecules may exist which could be characterized as MT-MDH proteins and which have as little as 30% identity over at least 150 residues to SEQ ID NO:1. The “95% variants” recited by the present claims have a variation that is far less than that of all potential MT-MDH proteins related to SEQ ID NO:1, i.e., those MT-MDH proteins having as little as 30% identity over at least 150 residues to SEQ ID NO:1. Therefore, one would expect the SEQ ID NO:1 variants recited by the present claims to have the functional activities of a MT-MDH protein.

As set forth in *In re Marzocchi*, 169 USPQ 367, 369 (CCPA 1971):

The first paragraph of § 112 ***requires nothing more than objective enablement.*** [emphasis added] How such a teaching is set forth, either by the use of illustrative examples or by broad terminology, is of no importance.

As a matter of Patent Office practice, then, a specification disclosure which contains a teaching of the manner and process of making and using the invention in terms which correspond in scope to those used in describing and defining the subject matter sought to be patented *must* be taken as in compliance with the enabling requirement of the first paragraph of § 112 *unless* there is reason to doubt the objective truth of the statements contained therein which must be relied on for enabling support.

Contrary to the standard set forth in *Marzocchi*, the Examiner has failed to provide any *reasons* why one would doubt that the guidance provided by the present Specification would enable one to make and use the recited variants of SEQ ID NO:1 or SEQ ID NO:2. Hence, a *prima facie* case for non-enablement has not been established with respect to the variants of SEQ ID NO:1 or SEQ ID NO:2.

For at least the above reasons, reversal of this rejection is requested.

(9) CONCLUSION

For at least the reasons set forth above, reversal of the rejections under 35 U.S.C. § 112, first paragraph, based on lack of an adequate written description and on lack of enablement, is respectfully requested.


If the USPTO determines that any additional fees are due, the Commissioner is hereby authorized to charge Deposit Account No. **09-0108**.

This brief is enclosed in triplicate.

Respectfully submitted,

INCYTE CORPORATION

Date: 06 January 2004




Richard C. Ekstrom

Reg. No. 37,027

Direct Dial Telephone: (650) 843-7352

Date: 1/06/04



Joel Harris

Reg. No. 44,743

Direct Dial Telephone: (650) 845-4866


3160 Porter Drive
Palo Alto, California 94304
Phone: (650) 855-0555
Fax: (650) 849-8886

Enclosures:

1. Brenner et al., Proc. Natl. Acad. Sci. U.S.A. 95:6073-78 (1998).

APPENDIX - CLAIMS ON APPEAL

3. An isolated polynucleotide encoding a polypeptide selected from the group consisting of:
- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1; and
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:1.
-
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. A cell transformed with a recombinant polynucleotide of claim 6.
9. A method for producing a polypeptide encoded by a polynucleotide of claim 3, the method comprising:
- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide of claim 3, and
 - b) recovering the polypeptide so expressed.
12. An isolated polynucleotide selected from the group consisting of:
- a) a polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2,
 - b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 95% identical to the polynucleotide sequence of SEQ ID NO:2,
 - c) a polynucleotide having a sequence complementary to a polynucleotide of a),
 - d) a polynucleotide having a sequence complementary to a polynucleotide of b) and
 - e) an RNA equivalent of a)-d).



Assessing sequence comparison methods with reliable structurally identified distant evolutionary relationships

STEVEN E. BRENNER^{*†‡}, CYRUS CHOTHIA^{*}, AND TIM J. P. HUBBARD[§]

^{*}MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, United Kingdom; and [†]Sanger Centre, Wellcome Trust Genome Campus, Hinxton, Cambs CB10 1SA, United Kingdom

Communicated by David R. Davies, National Institute of Diabetes, Bethesda, MD, March 16, 1998 (received for review November 12, 1997)

ABSTRACT Pairwise sequence comparison methods have been assessed using proteins whose relationships are known reliably from their structures and functions, as described in the SCOP database [Murzin, A. G., Brenner, S. E., Hubbard, T. & Chothia C. (1995) *J. Mol. Biol.* 247, 536–540]. The evaluation tested the programs BLAST [Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). *J. Mol. Biol.* 215, 403–410], WU-BLAST2 [Altschul, S. F. & Gish, W. (1996) *Methods Enzymol.* 266, 460–480], FASTA [Pearson, W. R. & Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. USA* 85, 2444–2448], and SSEARCH [Smith, T. F. & Waterman, M. S. (1981) *J. Mol. Biol.* 147, 195–197] and their scoring schemes. The error rate of all algorithms is greatly reduced by using statistical scores to evaluate matches rather than percentage identity or raw scores. The E-value statistical scores of SSEARCH and FASTA are reliable: the number of false positives found in our tests agrees well with the scores reported. However, the P-values reported by BLAST and WU-BLAST2 exaggerate significance by orders of magnitude. SSEARCH, FASTA $k_{\text{rup}} = 1$, and WU-BLAST2 perform best, and they are capable of detecting almost all relationships between proteins whose sequence identities are $>30\%$. For more distantly related proteins, they do much less well; only one-half of the relationships between proteins with 20–30% identity are found. Because many homologs have low sequence similarity, most distant relationships cannot be detected by any pairwise comparison method; however, those which are identified may be used with confidence.

Sequence database searching plays a role in virtually every branch of molecular biology and is crucial for interpreting the sequences issuing forth from genome projects. Given the method's central role, it is surprising that overall and relative capabilities of different procedures are largely unknown. It is difficult to verify algorithms on sample data because this requires large data sets of proteins whose evolutionary relationships are known unambiguously and independently of the methods being evaluated. However, nearly all known homologs have been identified by sequence analysis (the method to be tested). Also, it is generally very difficult to know, in the absence of structural data, whether two proteins that lack clear sequence similarity are unrelated. This has meant that although previous evaluations have helped improve sequence comparison, they have suffered from insufficient, imperfectly characterized, or artificial test data. Assessment also has been problematic because high quality database sequence searching attempts to have both sensitivity (detection of homologs) and specificity (rejection of unrelated proteins); however, these complementary goals are linked such that increasing one causes the other to be reduced.

Sequence comparison methodologies have evolved rapidly, so no previously published tests have evaluated modern versions of programs commonly used. For example, parameters in BLAST (1) have changed, and WU-BLAST2 (2)—which produces gapped alignments—has become available. The latest version of FASTA (3) previously tested was 1.6, but the current release (version 3.0) provides fundamentally different results in the form of statistical scoring.

The previous reports also have left gaps in our knowledge. For example, there has been no published assessment of thresholds for scoring schemes more sophisticated than percentage identity. Thus, the widely discussed statistical scoring measures have never actually been evaluated on large databases of real proteins. Moreover, the different scoring schemes commonly in use have not been compared.

Beyond these issues, there is a more fundamental question: in an absolute sense, how well does pairwise sequence comparison work? That is, what fraction of homologous proteins can be detected using modern database searching methods?

In this work, we attempt to answer these questions and to overcome both of the fundamental difficulties that have hindered assessment of sequence comparison methodologies. First, we use the set of distant evolutionary relationships in the SCOP: Structural Classification of Proteins database (4), which is derived from structural and functional characteristics (5). The SCOP database provides a uniquely reliable set of homologs, which are known independently of sequence comparison. Second, we use an assessment method that jointly measures both sensitivity and specificity. This method allows straightforward comparison of different sequence searching procedures. Further, it can be used to aid interpretation of real database searches and thus provide optimal and reliable results.

Previous Assessments of Sequence Comparison. Several previous studies have examined the relative performance of different sequence comparison methods. The most encompassing analyses have been by Pearson (6, 7), who compared the three most commonly used programs. Of these, the Smith–Waterman algorithm (8) implemented in SSEARCH (3) is the oldest and slowest but the most rigorous. Modern heuristics have provided BLAST (1) the speed and convenience to make it the most popular program. Intermediate between these two is FASTA (3), which may be run in two modes offering either greater speed ($k_{\text{rup}} = 2$) or greater effectiveness ($k_{\text{rup}} = 1$). Pearson also considered different parameters for each of these programs.

To test the methods, Pearson selected two representative proteins from each of 67 protein superfamilies defined by the PIR database (9). Each was used as a query to search the database, and the matched proteins were marked as being homologous or unrelated according to their membership of PIR.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/956073-6\$2.00/0
PNAS is available online at <http://www.pnas.org>.

Abbreviation: EPO, errors per query.

[†]Present address: Department of Structural Biology, Stanford University, Fairchild Building D-109, Stanford, CA 94305-5126

[‡]To whom reprints requests should be addressed. e-mail: brenner@hyper.stanford.edu.

superfamilies. Pearson found that modern matrices and "ln-scaling" of raw scores improve results considerably. He also reported that the rigorous Smith-Waterman algorithm worked slightly better than FASTA, which was in turn more effective than BLAST.

Very large scale analyses of matrices have been performed (10), and Henikoff and Henikoff (11) also evaluated the effectiveness of BLAST and FASTA. Their test with BLAST considered the ability to detect homologs above a predetermined score but had no penalty for methods which also reported large numbers of spurious matches. The Henikoffs searched the SWISS-PROT database (12) and used PROSITE (13) to define homologous families. Their results showed that the BLOSUM62 matrix (14) performed markedly better than the extrapolated PAM-series matrices (15), which previously had been popular.

A crucial aspect of any assessment is the data that are used to test the ability of the program to find homologs. But in Pearson's and the Henikoffs' evaluations of sequence comparison, the correct results were effectively unknown. This is because the superfamilies in PIR and PROSITE are principally created by using the same sequence comparison methods which are being evaluated. Interdependency of data and methods creates a "chicken and egg" problem, and means for example, that new methods would be penalized for correctly identifying homologs missed by older programs. For instance, immunoglobulin variable and constant domains are clearly homologous, but PIR places them in different superfamilies. The problem is widespread: each superfamily in PIR 48.00 with a structural homolog is itself homologous to an average of 1.6 other PIR superfamilies (16).

To surmount these sorts of difficulties, Sander and Schneider (17) used protein structures to evaluate sequence comparison. Rather than comparing different sequence comparison algorithms, their work focused on determining a length-dependent threshold of percentage identity, above which all proteins would be of similar structure. A result of this analysis was the HSSP equation; it states that proteins with 25% identity over 80 residues will have similar structures, whereas shorter alignments require higher identity. (Other studies also have used structures (18-20), but these focused on a small number of model proteins and were principally oriented toward evaluating alignment accuracy rather than homology detection.)

A general solution to the problem of scoring comes from statistical measures (i.e., E-values and P-values) based on the extreme value distribution (21). Extreme value scoring was implemented analytically in the BLAST program using the Karlin and Altschul statistics (22, 23) and empirical approaches have been recently added to FASTA and SSEARCH. In addition to being heralded as a reliable means of recognizing significantly similar proteins (24, 25), the mathematical tractability of statistical scores "is a crucial feature of the BLAST algorithm" (1). The validity of this scoring procedure has been tested analytically and empirically (see ref. 2 and references in ref. 24). However, all large empirical tests used random sequences that may lack the subtle structure found within biological sequences (26, 27) and obviously do not contain any real homologs. Thus, although many researchers have suggested that statistical scores be used to rank matches (24, 25, 28), there have been no large rigorous experiments on biological data to determine the degree to which such rankings are superior.

A Database for Testing Homology Detection. Since the discovery that the structures of hemoglobin and myoglobin are very similar though their sequences are not (29), it has been apparent that comparing structures is a more powerful (if less convenient) way to recognize distant evolutionary relationships than comparing sequences. If two proteins show a high degree of similarity in their structural details and function, it

is very probable that they have an evolutionary relationship though their sequence similarity may be low.

The recent growth of protein structure information combined with the comprehensive evolutionary classification in the SCOP database (4, 5) have allowed us to overcome previous limitations. With these data, we can evaluate the performance of sequence comparison methods on real protein sequences whose relationships are known confidently. The SCOP database uses structural information to recognize distant homologs, the large majority of which can be determined unambiguously. These superfamilies, such as the globins or the immunoglobulins, would be recognized as related by the vast majority of the biological community despite the lack of high sequence similarity.

From SCOP, we extracted the sequences of domains of proteins in the Protein Data Bank (PDB) (30) and created two databases. One (PDB90D-B) has domains which were all <90% identical to any other, whereas (PDB40D-B) had those <40% identical. The databases were created by first sorting all protein domains in SCOP by their quality and making a list. The highest quality domain was selected for inclusion in the database and removed from the list. Also removed from the list (and discarded) were all other domains above the threshold level of identity to the selected domain. This process was repeated until the list was empty. The PDB40D-B database contains 1,323 domains, which have 9,044 ordered pairs of distant relationships, or ~0.5% of the total 1,749,006 ordered pairs. In PDB90D-B, the 2,079 domains have 53,988 relationships, representing 1.2% of all pairs. Low complexity regions of sequence can achieve spurious high scores, so these were masked in both databases by processing with the SEG program (27) using recommended parameters: 12 1.8 2.0. The databases used in this paper are available from <http://sss.stanford.edu/sss/>, and databases derived from the current version of SCOP may be found at <http://scop.mrc-lmb.cam.ac.uk/scop/>.

Analyses from both databases were generally consistent, but PDB40D-B focuses on distantly related proteins and reduces the heavy overrepresentation in the PDB of a small number of families (31, 32), whereas PDB90D-B (with more sequences) improves evaluations of statistics. Except where noted otherwise, the distant homolog results here are from PDB40D-B. Although the precise numbers reported here are specific to the structural domain databases used, we expect the trends to be general.

Assessment Data and Procedure. Our assessment of sequence comparison may be divided into four different major categories of tests. First, using just a single sequence comparison algorithm at a time, we evaluated the effectiveness of different scoring schemes. Second, we assessed the reliability of scoring procedures, including an evaluation of the validity of statistical scoring. Third, we compared sequence comparison algorithms (using the optimal scoring scheme) to determine their relative performance. Fourth, we examined the distribution of homologs and considered the power of pairwise sequence comparison to recognize them. All of the analyses used the databases of structurally identified homologs and a new assessment criterion.

The analyses tested BLAST (1), version 1.4.9MP, and WU-BLAST2 (2), version 2.0a13MP. Also assessed was the FASTA package, version 3.0i76 (3), which provided FASTA and the SSEARCH implementation of Smith-Waterman (8). For SSEARCH and FASTA, we used BLOSUM45 with gap penalties -12/-1 (7, 16). The default parameters and matrix (BLOSUM62) were used for BLAST and WU-BLAST2.

The "Coverage Vs. Error" Plot. To test a particular protocol (comprising a program and scoring scheme), each sequence from the database was used as a query to search the database. This yielded ordered pairs of query and target sequences with associated scores, which were sorted, on the basis of their scores, from best to worst. The ideal method would have




FIG. 1. Coverage vs. error plots of different scoring schemes for SSEARCH Smith-Waterman. (A) Analysis of PDB400-B database. (B) Analysis of PDB900-B database. All of the proteins in the database were compared with each other using the SSEARCH program. The results of this single set of comparisons were considered using five different scoring schemes and assessed. The graphs show the coverage and errors per query (EPQ) for statistical scores, raw scores, and three measures using percentage identity. In the coverage vs. error plot, the x axis indicates the fraction of all homologs in the database (known from structure) which have been detected. Precisely, it is the number of detected pairs of proteins with the same fold divided by the total number of pairs from a common superfamily. PDB400-B contains a total of 9,044 homologs, so a score of 10% indicates identification of 904 relationships. The y axis reports the number of EPQ. Because there are 1,323 queries made in the PDB400-B all-vs.-all comparison, 13 errors corresponds to 0.01, or 1% EPQ. The y axis is presented on a log scale to show results over the widely varying degrees of accuracy which may be desired. The scores that correspond to the levels of EPQ and coverage are shown in Fig. 4 and Table 1. The graph demonstrates the trade-off between sensitivity and selectivity. As more homologs are found (moving to the right), more errors are made (moving up). The ideal method would be in the lower right corner of the graph, which corresponds to identifying many evolutionary relationships without selecting unrelated proteins. Three measures of percentage identity are plotted. Percentage identity within alignment is the degree of identity within the aligned region of the proteins, without consideration of the alignment length. Percentage identity within both is the number of identical residues in the aligned region as a percentage of the average length of the query and target proteins. The HSSP equation (17) is $H = 290.15 / (1 - 0.56^I)$ where I is length for $10 < I < 80$; $H > 100$ for $I < 10$; $H = 24.7$ for $I > 80$. The percentage identity HSSP-adjusted score is the percent identity within the alignment minus H . Smith-Waterman raw scores and E-values were taken directly from the sequence comparison program.

perfect separation, with all of the homologs at the top of the list and unrelated proteins below. In practice, perfect separation is impossible to achieve so instead one is interested in drawing a threshold above which there are the largest number of related pairs of sequences consistent with an acceptable error rate.

Our procedure involved measuring the coverage and error for every threshold. Coverage was defined as the fraction of structurally determined homologs that have scores above the selected threshold; this reflects the sensitivity of a method. Errors per query (EPQ), an indicator of selectivity, is the number of nonhomologous pairs above the threshold divided by the number of queries. Graphs of these data, called coverage vs. error plots, were devised to understand how

protocols compare at different levels of accuracy. These graphs share effectively all of the beneficial features of Receiver Operating Characteristic (ROC) plots (33, 34) but better represent the high degrees of accuracy required in sequence comparison and the huge background of nonhomologs.

This assessment procedure is directly relevant to practical sequence database searching, for it provides precisely the information necessary to perform a reliable sequence database search. The EPQ measure places a premium on score consistency; that is, it requires scores to be comparable for different queries. Consistency is an aspect which has been largely




FIG. 2. Unrelated proteins with high percentage identity. Hemoglobin β -chain (PDB code 1hds chain b, ref. 38, Left) and cellulase E2 (PDB code 1tml, ref. 39, Right) have 39% identity over 64 residues, a level which is often believed to be indicative of homology. Despite this high degree of identity, their structures strongly suggest that these proteins are not related. Appropriately, neither the raw alignment score of 85 nor the E-value of 1.3 is significant. Proteins rendered by RASMO (40).




FIG. 3. Length and percentage identity of alignments of unrelated proteins in PDB900-B: Each pair of nonhomologous proteins found with SSEARCH is plotted as a point whose position indicates the length and the percentage identity within the alignment. Because alignment length and percentage identity are quantized, many pairs of proteins may have exactly the same alignment length and percentage identity. The line shows the HSSP threshold (though it is intended to be applied with a different matrix and parameters).




FIG. 4. Reliability of statistical scores in PDB90D-B: Each line shows the relationship between reported statistical score and actual error rate for a different program. E-values are reported for SSEARCH and FASTA, whereas P-values are shown for BLAST and WU-BLAST2. If the scoring were perfect, then the number of errors per query and the E-values would be the same, as indicated by the upper bold line. (P-values should be the same as EPO for small numbers, and diverges at higher values, as indicated by the lower bold line.) E-values from SSEARCH and FASTA are shown to have good agreement with EPO but underestimate the significance slightly. BLAST and WU-BLAST2 are overconfident, with the degree of exaggeration dependent upon the score. The results for PDB40D-B were similar to those for PDB90D-B despite the difference in number of homologs detected. This graph could be used to roughly calibrate the reliability of a given statistical score.

ignored in previous tests but is essential for the straightforward or automatic interpretation of sequence comparison results. Further, it provides a clear indication of the confidence that should be ascribed to each match. Indeed, the EPQ measure should approximate the expectation value reported by database searching programs, if the programs' estimates are accurate.

The Performance of Scoring Schemes. All of the programs tested could provide three fundamental types of scores. The first score is the percentage identity, which may be computed in several ways based on either the length of the alignment or the lengths of the sequences. The second is a "raw" or "Smith-Waterman" score, which is the measure optimized by the Smith-Waterman algorithm and is computed by summing the substitution matrix scores for each position in the alignment and subtracting gap penalties. In BLAST, a measure





FIG. 5. Coverage vs. error plots of different sequence comparison methods: Five different sequence comparison methods are evaluated, each using statistical scores (E- or P-values). (A) PDB40D-B database. In this analysis, the best method is the slow SSEARCH, which finds 18% of relationships at 1% EPO. FASTA ktup = 1 and WU-BLAST2 are almost as good. (B) PDB90D-B database. The quick WU-BLAST2 program provides the best coverage at 1% EPO on this database, although at higher levels of error it becomes slightly worse than FASTA ktup = 1 and SSEARCH.

related to this score is scaled into bits. Third is a statistical score based on the extreme value distribution. These results are summarized in Fig. 1.

Sequence Identity. Though it has been long established that percentage identity is a poor measure (35), there is a common rule-of-thumb stating that 30% identity signifies homology. Moreover, publications have indicated that 25% identity can be used as a threshold (17, 36). We find that these thresholds, originally derived years ago, are not supported by present results. As databases have grown, so have the possibilities for chance alignments with high identity: thus, the reported cutoffs lead to frequent errors. Fig. 2 shows one of the many pairs of proteins with very different structures that nonetheless have high levels of identity over considerable aligned regions. Despite the high identity, the raw and the statistical scores for such incorrect matches are typically not significant. The principal reasons percentage identity does so poorly seem to be that it ignores information about gaps and about the conservative or radical nature of residue substitutions.

From the PDB90D-B analysis in Fig. 3, we learn that 30% identity is a reliable threshold for this database only for sequence alignments of at least 150 residues. Because one unrelated pair of proteins has 43.5% identity over 62 residues, it is probably necessary for alignments to be at least 70 residues in length before 40% is a reasonable threshold, for a database of this particular size and composition.

At a given reliability, scores based on percentage identity detect just a fraction of the distant homologs found by statistical scoring. If one measures the percentage identity in the aligned regions without consideration of alignment length, then a negligible number of distant homologs are detected. Use of the HSP equation improves the value of percentage identity, but even this measure can find only 4% of all known homologs at 1% EPQ. In short, percentage identity discards most of the information measured in a sequence comparison.

Raw Scores. Smith-Waterman raw scores perform better than percentage identity (Fig. 1), but ln-scaling (7) provided no notable benefit in our analysis. It is necessary to be very precise when using either raw or bit scores because a 20% change in cutoff score could yield a tenfold difference in EPQ. However, it is difficult to choose appropriate thresholds because the reliability of a bit score depends on the lengths of the proteins matched and the size of the database. Raw score thresholds also are affected by matrix and gap parameters.

Statistical Scores. Statistical scores were introduced partly to overcome the problems that arise from raw scores. This scoring scheme provides the best discrimination between homologous proteins and those which are unrelated. Most

likely, its power can be attributed to its incorporation of more information than any other measure; it takes account of the full substitution and gap data (like raw scores) but also has details about the sequence lengths and composition and is scaled appropriately.

We find that statistical scores are not only powerful, but also easy to interpret. SSEARCH and FASTA show close agreement between statistical scores and actual number of errors per query (Fig. 4). The expectation value score gives a good, slightly conservative estimate of the chances of the two sequences being found at random in a given query. Thus, an E-value of 10.01 indicates that roughly one pair of nonhomologs of this similarity should be found in every 100 different queries. Neither raw scores nor percentage identity can be interpreted in this way, and these results validate the suitability of the extreme value distribution for describing the scores from a database search.

The P-values from BLAST also should be directly interpretable but were found to overstate significance by more than two orders of magnitude for 1% EPO for this database. Nonetheless, these results strongly suggest that the analytic theory is fundamentally appropriate. WU-BLAST2 scores were more reliable than those from BLAST, but also exaggerate expected confidence by more than an order of magnitude at 1% EPO.

Overall Detection of Homologs and Comparison of Algorithms. The results in Fig. 5A and Table 1 show that pairwise sequence comparison is capable of identifying only a small fraction of the homologous pairs of sequences in PDB40D-B. Even SSEARCH with E-values, the best protocol tested, could find only 18% of all relationships at a 1% EPO. BLAST, which identifies 15%, was the worst performer, whereas FASTA $k_{\text{up}} = 1$ is nearly as effective as SSEARCH. FASTA $k_{\text{up}} = 2$ and WU-BLAST2 are intermediate in their ability to detect homologs. Comparison of different algorithms indicates that those capable of identifying more homologs are generally slower. SSEARCH is 25 times slower than BLAST and 6.5 times slower than FASTA $k_{\text{up}} = 1$. WU-BLAST2 is slightly faster than FASTA $k_{\text{up}} = 2$, but the latter has more interpretable scores.

In PDB40D-B, where there are many close relationships, the best method can identify only 38% of structurally known homologs (Fig. 5B). The method which finds that many relationships is WU-BLAST2. Consequently, we infer that the differences between FASTA $k_{\text{up}} = 1$, SSEARCH, and WU-BLAST2 programs are unlikely to be significant when compared with variation in database composition and scoring reliability.

Fig. 6 helps to explain why most distant homologs cannot be found by sequence comparison: a great many such relationships have no more sequence identity than would be expected by chance. SSEARCH with E-values can recognize >90% of the homologous pairs with 30–40% identity. In this region, there are 30 pairs of homologous proteins that do not have significant E-values, but 26 of these involve sequences with <50 residues. Of sequences having 25–30% identity, 75% are identified by SSEARCH E-values. However, although the number of homologs grows at lower levels of identity, the detection falls off sharply: only 40% of homologs with 20–25% identity




FIG. 6. Distribution and detection of homologs in PDB40D-B. Bars show the distribution of homologous pairs PDB40D-B according to their identity (using the measure of identity in both). Filled regions indicate the number of these pairs found by the best database searching method (SSEARCH with E-values) at 1% EPO. The PDB40D-B database contains proteins with <40% identity, and as shown on this graph, most structurally identified homologs in the database have diverged extremely far in sequence and have <20% identity. Note that the alignments may be inaccurate, especially at low levels of identity. Filled regions show that SSEARCH can identify most relationships that have 25% or more identity, but its detection wanes sharply below 25%. Consequently, the great sequence divergence of most structurally identified evolutionary relationships effectively defeats the ability of pairwise sequence comparison to detect them.

are detected and only 10% of those with 15–20% can be found. These results show that statistical scores can find related proteins whose identity is remarkably low; however, the power of the method is restricted by the great divergence of many protein sequences.

After completion of this work, a new version of pairwise BLAST was released: BLASTGP (37). It supports gapped alignments, like WU-BLAST2, and dispenses with sum statistics. Our initial tests on BLASTGP using default parameters show that its E-values are reliable and that its overall detection of homologs was substantially better than that of ungapped BLAST, but not quite equal to that of WU-BLAST2.

CONCLUSION

The general consensus amongst experts (see refs. 7, 24, 25, 27 and references therein) suggests that the most effective sequence searches are made by (i) using a large current database in which the protein sequences have been complexity masked and (ii) using statistical scores to interpret the results. Our experiments fully support this view.

Our results also suggest two further points. First, the E-values reported by FASTA and SSEARCH give fairly accurate estimates of the significance of each match, but the P-values provided by BLAST and WU-BLAST2 underestimate the true

Table 1. Summary of sequence comparison methods with PDB40D-B

Method	Relative Time*	1% EPO Cutoff	Coverage at 1% EPO
SSEARCH % identity: within alignment	25.5	>70%	<0.1
SSEARCH % identity: within both	25.5	34%	3.0
SSEARCH % identity: HSSP-scaled	25.5	35% (HSSP = 9.8)	4.0
SSEARCH Smith-Waterman raw scores	25.5	142	10.5
SSEARCH E-values	25.5	0.03	18.4
FASTA $k_{\text{up}} = 1$ E-values	3.9	0.03	17.9
FASTA $k_{\text{up}} = 2$ E-values	1.4	0.03	16.7
WU-BLAST2 P-values	1.1	0.003	17.5
BLAST P-values	1.0	0.00016	14.8

*Times are from large database searches with genome proteins.

extent of errors. Second, SSEARCH, WU-BLAST2, and FASTA ktup = 1 perform best, though BLAST and FASTA ktup = 2 detect most of the relationships found by the best procedures and are appropriate for rapid initial searches.

The homologous proteins that are found by sequence comparison can be distinguished with high reliability from the huge number of unrelated pairs. However, even the best database searching procedures tested fail to find the large majority of distant evolutionary relationships at an acceptable error rate. Thus, if the procedures assessed here fail to find a reliable match, it does not imply that the sequence is unique; rather, it indicates that any relatives it might have are distant ones.**

**Additional and updated information about this work, including supplementary figures, may be found at <http://sss.stanford.edu/sss/>.

The authors are grateful to Drs. A. G. Murzin, M. Levitt, S. R. Eddy, and G. Mitchison for valuable discussion. S.E.B. was principally supported by a St. John's College (Cambridge, UK) Benefactors' Scholarship and by the American Friends of Cambridge University. S.E.B. dedicates his contribution to the memory of Rabbi Albert T. and Clara S. Bilgray.

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403-410.
- Altschul, S. F. & Gish, W. (1996) *Methods Enzymol.* 266, 460-480.
- Pearson, W. R. & Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. USA* 85, 2444-2448.
- Murzin, A. G., Brenner, S. E., Hubbard, T. & Chothia, C. (1995) *J. Mol. Biol.* 247, 536-540.
- Brenner, S. E., Chothia, C., Hubbard, T. J. P. & Murzin, A. G. (1996) *Methods Enzymol.* 266, 635-643.
- Pearson, W. R. (1991) *Genomics* 11, 635-650.
- Pearson, W. R. (1995) *Protein Sci.* 4, 1145-1160.
- Smith, T. F. & Waterman, M. S. (1981) *J. Mol. Biol.* 147, 195-197.
- George, D. G., Hunt, L. T. & Barker, W. C. (1996) *Methods Enzymol.* 266, 41-59.
- Vogt, G., Etzold, T. & Argos, P. (1995) *J. Mol. Biol.* 249, 816-831.
- Henikoff, S. & Henikoff, J. G. (1993) *Proteins* 17, 49-61.
- Bairoch, A. & Apweiler, R. (1996) *Nucleic Acids Res.* 24, 21-25.
- Bairoch, A., Bucher, P. & Hofmann, K. (1996) *Nucleic Acids Res.* 24, 189-196.
- Henikoff, S. & Henikoff, J. G. (1992) *Proc. Natl. Acad. Sci. USA* 89, 10915-10919.
- Dayhoff, M., Schwartz, R. M. & Orcutt, B. C. (1978) in *Atlas of Protein Sequence and Structure*, ed. Dayhoff, M. (National Bio-medical Research Foundation, Silver Spring, MD), Vol. 5, Suppl. 3, pp. 345-352.
- Brenner, S. E. (1996) Ph.D. thesis (University of Cambridge, UK).
- Sander, C. & Schneider, R. (1991) *Proteins* 9, 56-68.
- Johnson, M. S. & Overington, J. P. (1993) *J. Mol. Biol.* 233, 716-738.
- Barton, G. J. & Sternberg, M. J. E. (1987) *Protein Eng.* 1, 89-94.
- Lesk, A. M., Levitt, M. & Chothia, C. (1986) *Protein Eng.* 1, 77-78.
- Arratia, R., Gordon, L. & M. W. (1986) *Ann. Stat.* 14, 971-993.
- Karlin, S. & Altschul, S. F. (1990) *Proc. Natl. Acad. Sci. USA* 87, 2264-2268.
- Karlin, S. & Altschul, S. F. (1993) *Proc. Natl. Acad. Sci. USA* 90, 5873-5877.
- Altschul, S. F., Boguski, M. S., Gish, W. & Wootton, J. C. (1994) *Nat. Genet.* 6, 119-129.
- Pearson, W. R. (1996) *Methods Enzymol.* 266, 227-258.
- Lipman, D. J., Wilbur, W. J., Smith, T. F. & Waterman, M. S. (1984) *Nucleic Acids Res.* 12, 215-226.
- Wootton, J. C. & Federhen, S. (1996) *Methods Enzymol.* 266, 554-571.
- Waterman, M. S. & Vingron, M. (1994) *Stat. Science* 9, 367-381.
- Perutz, M. F., Kendrew, J. C. & Watson, H. C. (1965) *J. Mol. Biol.* 13, 669-678.
- Abola, E. E., Bernstein, F. C., Bryant, S. H., Koetzle, T. F. & Weng, J. (1987) in *Crystallographic Databases: Information Content, Software Systems, Scientific Applications*, eds. Allen, F. H., Bergerhoff, G. & Sievers, R. (Data Comm. Intl. Union Crystallogr., Cambridge, UK), pp. 107-132.
- Brenner, S. E., Chothia, C. & Hubbard, T. J. P. (1997) *Curr. Opin. Struct. Biol.* 7, 369-376.
- Orengo, C., Michie, A., Jones, S., Jones, D. T., Swindells, M. B. & Thornton, J. (1997) *Structure (London)* 5, 1093-1108.
- Zweig, M. H. & Campbell, G. (1993) *Clin. Chem.* 39, 561-577.
- Gribskov, M. & Robinson, N. L. (1996) *Comput. Chem.* 20, 25-33.
- Fitch, W. M. (1966) *J. Mol. Biol.* 16, 9-16.
- Chung, S. Y. & Subbiah, S. (1996) *Structure (London)* 4, 1123-1127.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997) *Nucleic Acids Res.* 25, 3389-3402.
- Girling, R., Schmidt, W., Jr., Houston, T., Amma, E. & Huisman, T. (1979) *J. Mol. Biol.* 131, 417-433.
- Spezio, M., Wilson, D. & Karplus, P. (1993) *Biochemistry* 32, 9906-9916.
- Sayle, R. A. & Milner-White, E. J. (1995) *Trends Biochem. Sci.* 20, 374-376.